

GLUEING OPERATION FOR R-MATRICES, QUANTUM GROUPS AND LINK-INVARIANTS OF HECKE TYPE

Shahn MAJID¹

Department of Applied Mathematics & Theoretical Physics
University of Cambridge, CB3 9EW, U.K.
e-mail: majid@damtp.cambridge.ac.uk

Martin MARKL

Mathematical Institute of the Academy
Žitná 25, 115 67 Prague, Czech Republic
e-mail: markl@csearn.bitnet

July 1993

Abstract. We introduce an associative glueing operation $\oplus_q$ on the space of solutions of the Quantum Yang-Baxter Equations of Hecke type. The corresponding glueing operations for the associated quantum groups and quantum vector spaces are also found. The former involves 2×2 quantum matrices whose entries are themselves square or rectangular quantum matrices. The corresponding glueing operation for link-invariants is introduced and involves a state-sum model with Boltzmann weights determined by the link invariants to be glued. The standard $su(n)$ solution, its associated quantum matrix group, quantum space and link-invariant arise at once by repeated glueing of the one-dimensional case.

1. Introduction.

Matrix representations of the Artin braid group, also called Yang-Baxter operators or R-matrices, have been extensively studied in the last several years. Each solution leads to interesting algebraic structures such as quantum groups [3] and braided groups [16][13] as well as (in the nice cases) to link-invariants. The celebrated Jones polynomial for knots and links is of this type and corresponds to the R-matrix for the quantum group $GL_q(2)$.

¹SERC Fellow and Fellow of Pembroke College, Cambridge

In spite of this great activity, the problem still remains how to obtain such Yang-Baxter operators. In concrete terms they are matrix solutions $R \in M_n \otimes M_n$ for some n of the cubic equations

$$(1) \quad R_{12}R_{13}R_{23} = R_{23}R_{13}R_{12}.$$

Here M_n denotes $n \times n$ matrices and $R_{12} = R \otimes 1$ in $M_n^{\otimes 3}$ etc. These cubic equations are highly-overdetermined and yet have a rich variety of solutions. In general, little is known about the moduli space of solutions for $n > 2$.

We study in this paper part of this moduli space, namely the solutions of (1) of Hecke type. These are solutions with quadratic minimal polynomials for the associated braid representation $\mathcal{R}$. Without loss of generality we can suppose that the minimal polynomial takes the form

$$(2) \quad (\mathcal{R} - q)(\mathcal{R} + q^{-1}) = 0$$

for some q which we fix. On the other hand we do not fix any size n , so we consider all Hecke solutions together of any dimension or infinite dimensional. Note that Hecke solutions have been studied in [6] but with other results than those here. The novel result in this paper is an associative glueing operation $\oplus_q$ for such solutions. This construction has a great many consequences. First, the standard solutions associated to the Lie algebra $su(n)$ (which are known to be of Hecke type) are understood now as the n -fold $\oplus_q$ of the trivial 1-dimensional solution $R = (q)$. Thus we answer the question how to obtain new R-matrices from old ones in a way that gives the standard non-trivial R-matrices from something trivial. This is the main result of Section 2, where formal definitions (in an operator notation) can be found. We also express our $\oplus_q$ operation in categorical terms in this section as some kind of direct sum.

Many of the constructions that are usually made with R -matrices or Yang-Baxter operators behave well under our glueing operation. Thus the quantum matrix bialgebra $M_q(n)$ (whose localization is the quantum group $GL_q(n)$) is built up in a natural way from an iterated glueing operation $*_q$. Since the construction is associative one has just as well

$$M_q(m+n) = M_q(m) *_q M_q(n), \quad \forall m, n \geq 1.$$

Equally well we can glue the quantum matrices $A(R)$ coming from any R-matrices of Hecke type to obtain new ones. This is the main result of Section 3. The general structure of the glueing $A(R) *_q A(R')$ here involves the novel idea of a 2×2 quantum block matrix whose entries are themselves quantum matrices. The off-diagonal blocks here involve the notion of an $m \times n$ quantum matrix algebra $A(R : R')$, which we also introduce.

In Section 4 we give the same construction of the associated quantum spaces or exchange algebras. This setting is the original motivation behind our construction. Namely, the natural structure behind these quantum spaces (such as the celebrated quantum plane $yx = qxy$) is that of a braided-vector-space. I.e., they form Hopf algebras in braided categories [13] with the braided-coproduct expressing linear addition of vectors. The quantum plane lives in the category associated to the $GL_q(2)$ R-matrix. On the other hand, there is no canonical tensor product for Hopf algebras in strictly braided categories because the usual tensor product construction becomes tangled up in the braided case. Our result provides a partial solution to this problem. At least in the Hecke case the braided vector space algebras can be tensor produced in some way provided their associated braidings are also glued according to our $\oplus_q$ construction. Thus

$$\mathbb{C}_q^{m+n} = \mathbb{C}_q^m \otimes_q \mathbb{C}_q^n, \quad \forall m, n \geq 1$$

as Hopf algebras in three different braided categories. Equally well we can glue quantum spaces associated to any two Hecke R-matrices to obtain a third one. The tensor product $\otimes_q$ needed here is the braided tensor product of $\mathbb{Z}$ -graded algebras along the lines introduced in [14].

We turn in Section 5 to the behaviour of quantum traces and ribbon elements under our $\oplus_q$ construction. These are needed for the construction of link-invariants as well as (much the same thing) to the construction of braided categories with duals associated to R-matrices. Some basic facts here about such dualisable R-matrices are developed in this section also.

Finally in Section 6 we conclude with a simple state-sum model description of the glueing of link-invariants of Hecke type to obtain new ones. The construction generalizes Kauffman's state model for the Jones polynomial. We do not expect to obtain genuinely new invariants (they should all be restrictions of the HOMFLY polynomial) but we do obtain a new associative operation among them.

2. Glueing of Yang-Baxter Operators.

All objects in this paper are assumed to be defined over a fixed field $\mathbf{k}$ of characteristic zero. Typically $\mathbf{k} = \mathbb{C}$, the field of complex numbers. As a suitable reference for the conventions we recommend [19].

Definition 2.1. *Let V be a vector space. By a Yang-Baxter operator on V we mean an invertible operator $\mathcal{R} \in \text{End}(V \otimes V)$ satisfying the following Quantum Yang-Baxter equation (QYBE):*

$$(\mathcal{R} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R})(\mathcal{R} \otimes \mathbf{1}) = (\mathbf{1} \otimes \mathcal{R})(\mathcal{R} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R}).$$

If V is finite dimensional with a basis (x_a) , we may introduce the matrix $R = (R^a_i{}^b_j)$ by

$$\mathcal{R}(x_i \otimes x_j) = x_b \otimes x_a R^a_i{}^b_j.$$

The QYBE written in terms of R then reads as in (1). This is the form familiar in physics so we shall state our results in this form as well as in the operator form with which we work.

Definition 2.2. *Let $q \in \mathbf{k}$ be an invertible element. By a q -Hecke operator we mean an element $\mathcal{R} \in \text{End}(V \otimes V)$ satisfying*

$$(\mathcal{R} - q)(\mathcal{R} + q^{-1}) = 0, \text{ or, equivalently, } \mathcal{R}^2 - (q - q^{-1})\mathcal{R} - \mathbf{1} = 0.$$

Definition 2.3. *1. Denote by $\mathcal{YB}$ the category whose objects are couples $(V, \mathcal{R})$, where V is a vector space and $\mathcal{R}$ is a Yang-Baxter operator on V . Morphisms from $(V, \mathcal{R})$ to $(W, \mathcal{S})$ in $\mathcal{YB}$ are linear maps $f : V \rightarrow W$ such that $\mathcal{S} \circ (f \otimes f) = (f \otimes f) \circ \mathcal{R}$.*

2. Denote by $\mathcal{H}_q$ the category whose objects are the couples $(V, \mathcal{R})$, where $\mathcal{R}$ is an Yang-Baxter q -Hecke operator on V and whose morphisms from $(V, \mathcal{R})$ to $(W, \mathcal{S})$ are linear maps $h : V \otimes V \rightarrow W \otimes W$ such that $\mathcal{S} \circ h = h \circ \mathcal{R}$.

Notice the substantial difference of the nature of morphisms in $\mathcal{YB}$ and $\mathcal{H}_q$. We will see the need for this later in the section.

One of the basic constructions of this paper is described in the following theorem.

Theorem 2.4. *Let $(X, \mathcal{R})$ and $(Y, \mathcal{R}')$ be elements of $\mathcal{H}_q$. Define their glueing $\mathcal{R} \oplus_q \mathcal{R}' \in \text{End}((X \oplus Y) \otimes (X \oplus Y))$ by*

$$\begin{aligned} (\mathcal{R} \oplus_q \mathcal{R}')|_{X \otimes X} &= \mathcal{R}, & (\mathcal{R} \oplus_q \mathcal{R}')|_{Y \otimes Y} &= \mathcal{R}', \\ (\mathcal{R} \oplus_q \mathcal{R}')(x \otimes y) &= y \otimes x, & (\mathcal{R} \oplus_q \mathcal{R}')(y \otimes x) &= x \otimes y + (q - q^{-1}) \cdot y \otimes x \end{aligned}$$

for $x \in X$ and $y \in Y$. Then $(X \oplus Y, \mathcal{R} \oplus_q \mathcal{R}') \in \mathcal{H}_q$.

Proof. We postpone the proof to 2.12 where we show that $\mathcal{R} \oplus_q \mathcal{R}'$ is a special case of a much more general construction described in Theorem 2.7. ■

Under a suitable choice of conventions, the corresponding matrix $R \oplus_q R' = ((R \oplus_q R')_{ij}^{ab})$ can be written in a nice block form. To be more precise, let $X = \text{Span}(x_k; 1 \leq k \leq m)$ and

$Y = \text{Span}(y_l; m+1 \leq k \leq m+n)$. If $I = \{kl; 1 \leq k, l \leq m+n\}$, then we can write $I = I_X \cup I_< \cup I_> \cup I_Y$ (the disjoint union), where

$$I_X = \{kl \in I; k, l \leq m\}, \quad I_Y = \{kl \in I; k, l \geq m+1\},$$

$$I_< = \{kl \in I; k \leq m \text{ \& } l \geq m+1\}, \quad I_> = \{kl \in I; k \geq m+1 \text{ \& } l \leq m\}.$$

We can order I by requiring that $I_X < I_< < I_> < I_Y$ and that each of the four pieces $I_X, I_Y, I_<$ and $I_>$ has the standard lexicographic order. Then

$$(3) \quad R \oplus_q R' = \begin{pmatrix} R & 0 & 0 & 0 \\ 0 & \mathbf{1} & (q - q^{-1})P & 0 \\ 0 & 0 & \mathbf{1} & 0 \\ 0 & 0 & 0 & R' \end{pmatrix}.$$

Here $(R \oplus_q R')_{ij}^{ab}$ is given as the above matrix entry at row ab and column ij in the order introduced above, where R (resp. R') is the matrix representing $\mathcal{R}$ (resp. $\mathcal{R}'$), $\mathbf{1}$ is the identity matrix and P is the permutation matrix, $P_{ij}^{ab} = \delta_j^a \delta_i^b$ (Kronecker deltas).

Example 2.5. Let $n \geq 1$ and let E_{ij} be, for $1 \leq i, j \leq n$, the $(n \times n)$ -matrix given by $(E_{ij})_l^k = \delta_i^k \delta_l^j$. Then the well-known Yang-Baxter operator $\mathcal{R}_{su(n)}$ related with the semisimple Lie algebra $su(n)$ and the matrix quantum groups $GL_q(n)$ and $SL_q(n)$ is given by the matrix

$$R_{su(n)} := (q \sum_i E_{ii} \otimes E_{ii} \oplus \sum_{i \neq j} E_{ii} \otimes E_{jj} + (q - q^{-1}) \sum_{j > i} E_{ij} \otimes E_{ji})$$

It can easily be seen that the corresponding operator is a glueing

$$\mathcal{R}_{su(m+n)} = \mathcal{R}_{su(m)} \oplus_q \mathcal{R}_{su(n)}, \text{ for } m, n \geq 1.$$

Thus the glueing enables us to construct $\mathcal{R}_{su(m)}$ inductively by

$$\mathcal{R}_{su(m+1)} = \mathcal{R}_{su(m)} \oplus_q \mathcal{R}_{su(1)}$$

from the one-dimensional solution $\mathcal{R}_{su(1)}(1 \otimes 1) = q1 \otimes 1$. This description of $\mathcal{R}_{su(n)}$ ensures that it possesses the q -Hecke property, as well as any other properties which hold for $\mathcal{R}_{su(1)}$ and which are preserved under $\oplus_q$.

Example 2.6. Let $\mathcal{R}, \mathcal{R}' \in \text{End}(\mathbf{k} \otimes \mathbf{k})$ be the Yang-Baxter q -Hecke operators given by $\mathcal{R}(1 \otimes 1) = q \cdot (1 \otimes 1)$ and $\mathcal{R}'(1 \otimes 1) = -q^{-1} \cdot (1 \otimes 1)$ respectively. Then $\mathcal{R} \oplus_q \mathcal{R}'$ is the non-standard Yang-Baxter q -Hecke operator $\mathcal{R}_{gl(1|1)}$ given by the matrix

$$\begin{pmatrix} q & 0 & 0 & 0 \\ 0 & 1 & q - q^{-1} & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -q^{-1} \end{pmatrix}$$

The glueing is a special case of a more general construction described in the following theorem.

Theorem 2.7. *Let X and Y be vector spaces and let $\mathcal{R} : X \otimes X \rightarrow X \otimes X$, $\mathcal{R}' : Y \otimes Y \rightarrow Y \otimes Y$, $\mathcal{U} : X \otimes Y \rightarrow Y \otimes X$, $\mathcal{S} : Y \otimes X \rightarrow X \otimes Y$ and $\mathcal{T} : Y \otimes X \rightarrow Y \otimes X$ be linear maps. Define $\mathcal{R} \oplus_{\mathcal{U}, \mathcal{S}, \mathcal{T}} \mathcal{R}' \in \text{End}((X \oplus Y) \otimes (X \oplus Y))$ by*

$$(\mathcal{R} \oplus_{\mathcal{U}, \mathcal{S}, \mathcal{T}} \mathcal{R}')|_{X \otimes X} = \mathcal{R}, \quad (\mathcal{R} \oplus_{\mathcal{U}, \mathcal{S}, \mathcal{T}} \mathcal{R}')|_{Y \otimes Y} = \mathcal{R}',$$

$$(\mathcal{R} \oplus_{\mathcal{U}, \mathcal{S}, \mathcal{T}} \mathcal{R}')(x \otimes y) = \mathcal{U}(x \otimes y), \quad (\mathcal{R} \oplus_{\mathcal{U}, \mathcal{S}, \mathcal{T}} \mathcal{R}')(y \otimes x) = \mathcal{S}(y \otimes x) + \mathcal{T}(y \otimes x),$$

for $x \in X$ and $y \in Y$. This defines a Yang-Baxter operator on $X \oplus Y$ if and only if both $\mathcal{R}$ and $\mathcal{R}'$ are Yang-Baxter operators, $\mathcal{U}$ and $\mathcal{S}$ are invertible, and the following conditions are satisfied

$$(I) \quad (\mathbf{1} \otimes \mathcal{R})(\mathcal{U} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{U}) = (\mathcal{U} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{U})(\mathcal{R} \otimes \mathbf{1}) \text{ on } X^{\otimes 2} \otimes Y,$$

$$(I)' \quad (\mathcal{R}' \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{U})(\mathcal{U} \otimes \mathbf{1}) = (\mathbf{1} \otimes \mathcal{U})(\mathcal{U} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R}') \text{ on } X \otimes Y^{\otimes 2},$$

$$(II) \quad (\mathbf{1} \otimes \mathcal{S})(\mathcal{S} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R}) = (\mathcal{R} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{S})(\mathcal{S} \otimes \mathbf{1}) \text{ on } Y \otimes X^{\otimes 2},$$

$$(II)' \quad (\mathcal{S} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{S})(\mathcal{R}' \otimes \mathbf{1}) = (\mathbf{1} \otimes \mathcal{R}')(\mathcal{S} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{S}) \text{ on } Y^{\otimes 2} \otimes X,$$

$$(III) \quad (\mathbf{1} \otimes \mathcal{U})(\mathcal{R} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{S}) = (\mathcal{S} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R})(\mathcal{U} \otimes \mathbf{1}) \text{ on } X \otimes Y \otimes X,$$

$$(III)' \quad (\mathcal{U} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R}')(\mathcal{S} \otimes \mathbf{1}) = (\mathbf{1} \otimes \mathcal{S})(\mathcal{R}' \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{U}) \text{ on } Y \otimes X \otimes Y,$$

$$(IV) \quad (\mathbf{1} \otimes \mathcal{R})(\mathcal{U} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{T}) = (\mathcal{T} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R})(\mathcal{U} \otimes \mathbf{1}) \text{ on } X \otimes Y \otimes X,$$

$$(IV)' \quad (\mathcal{R}' \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{U})(\mathcal{T} \otimes \mathbf{1}) = (\mathbf{1} \otimes \mathcal{T})(\mathcal{R}' \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{U}) \text{ on } Y \otimes X \otimes Y,$$

$$(V) \quad (\mathbf{1} \otimes \mathcal{T})(\mathcal{S} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R}) = (\mathcal{S} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R})(\mathcal{T} \otimes \mathbf{1}) \text{ on } Y \otimes X^{\otimes 2},$$

$$(V)' \quad (\mathcal{T} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{S})(\mathcal{R}' \otimes \mathbf{1}) = (\mathbf{1} \otimes \mathcal{S})(\mathcal{R}' \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{T}) \text{ on } X \otimes Y^{\otimes 2},$$

$$(VI) \quad (\mathbf{1} \otimes \mathcal{R})(\mathcal{T} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R}) = (\mathcal{T} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R})(\mathcal{T} \otimes \mathbf{1}) + (\mathcal{U} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{T})(\mathcal{S} \otimes \mathbf{1}) \text{ on } Y \otimes X^{\otimes 2},$$

$$(VI)' \quad (\mathcal{R}' \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{T})(\mathcal{R}' \otimes \mathbf{1}) = (\mathbf{1} \otimes \mathcal{T})(\mathcal{R}' \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{T}) + (\mathbf{1} \otimes \mathcal{U})(\mathcal{T} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{S}) \text{ on } X \otimes Y^{\otimes 2}.$$

Proof: First, it is immediate to see that $\mathcal{R} \oplus_{\mathcal{U}, \mathcal{S}, \mathcal{T}} \mathcal{R}'$ is invertible if and only if both $\mathcal{R}$, $\mathcal{R}'$, $\mathcal{S}$ and $\mathcal{U}$ are invertible operators.

We shall verify that $\mathcal{Q} = \mathcal{R} \oplus_{\mathcal{U}, \mathcal{S}, \mathcal{T}} \mathcal{R}'$ satisfies the QYBE on $(X \oplus Y)^{\otimes 3}$, which can be decomposed as

$$(4) \ X^{\otimes 3} \oplus (Y \otimes X^{\otimes 2}) \oplus (X \otimes Y \otimes X) \oplus (X^{\otimes 2} \otimes Y) \oplus (X \otimes Y^{\otimes 2}) \oplus (Y \otimes X \otimes Y) \oplus (Y^{\otimes 2} \otimes X) \oplus Y^{\otimes 3}.$$

On the first component of the previous decomposition we clearly have

$$(\mathcal{Q} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{Q})(\mathcal{Q} \otimes \mathbf{1})|_{X^{\otimes 3}} = (\mathcal{R} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R})(\mathcal{R} \otimes \mathbf{1})$$

and, similarly,

$$(\mathbf{1} \otimes \mathcal{Q})(\mathcal{Q} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{Q})|_{X^{\otimes 3}} = (\mathbf{1} \otimes \mathcal{R})(\mathcal{R} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R}),$$

therefore QYBE for $\mathcal{Q}$ on the first component of (4) is equivalent with QYBE for $\mathcal{R}$. On the next component of (4) we have

$$\begin{aligned} (\mathcal{Q} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{Q})(\mathcal{Q} \otimes \mathbf{1})|_{(Y \otimes X^{\otimes 2})} &= (\mathcal{R} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{S})(\mathcal{S} \otimes \mathbf{1}) + \\ &+ (\mathcal{U} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{T})(\mathcal{S} \otimes \mathbf{1}) + (\mathcal{S} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R})(\mathcal{T} \otimes \mathbf{1}) + (\mathcal{T} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R})(\mathcal{T} \otimes \mathbf{1}), \end{aligned}$$

while

$$\begin{aligned} (\mathbf{1} \otimes \mathcal{Q})(\mathcal{Q} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{Q})|_{(Y \otimes X^{\otimes 2})} &= \\ &= (\mathbf{1} \otimes \mathcal{S})(\mathcal{S} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R}) + (\mathbf{1} \otimes \mathcal{T})(\mathcal{S} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R}) + (\mathbf{1} \otimes \mathcal{R})(\mathcal{T} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R}). \end{aligned}$$

The terms of the above two equations are maps from $(Y \otimes X^{\otimes 2})$ to $(Y \otimes X^{\otimes 2}) \oplus (X \otimes Y \otimes X) \oplus (X^{\otimes 2} \otimes Y)$. Looking at the component in $(Y \otimes X^{\otimes 2})$ we get that the QYBE for $\mathcal{Q}$ imply

$$(\mathcal{U} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{T})(\mathcal{S} \otimes \mathbf{1}) + (\mathcal{T} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R})(\mathcal{T} \otimes \mathbf{1}) = (\mathbf{1} \otimes \mathcal{R})(\mathcal{T} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{R}),$$

which is (VI). Similarly, the component in $(X \otimes Y \otimes X)$ gives (V) and the component in $(X^{\otimes 2} \otimes Y)$ gives (II). Repeating the same discussion for the remaining six components of (4) we get all the equations (I)–(VI)' of the theorem. On the other hand, it also follows immediately from this discussion that the equations (I)–(VI)' imply the QYBE for $\mathcal{Q}$. ■

This provides an extension of our glueing construction to a more general situation (and is not directly connected with the Hecke situation). Using the obvious similar conventions as in (3),

we can represent the resulting Yang-Baxter operator by the matrix

$$(5) \quad R \oplus_{U,S,T} R' = \begin{pmatrix} R & 0 & 0 & 0 \\ 0 & U & T & 0 \\ 0 & 0 & S & 0 \\ 0 & 0 & 0 & R' \end{pmatrix}$$

where the matrices R, R', U, S, T are matrices representing the operators in the theorem. In matrix terms the conditions (I)-(VI) in theorem are

$$U_{12}U_{13}R_{23} = R_{23}U_{13}U_{12}, \quad R_{12}S_{13}S_{23} = S_{23}S_{13}R_{12}, \quad S_{12}R_{13}U_{23} = U_{23}R_{13}S_{12}$$

$$T_{12}U_{13}R_{23} = U_{23}R_{13}T_{12}, \quad R_{12}S_{13}T_{23} = S_{23}R_{13}T_{12}, \quad R_{12}T_{13}R_{23} = T_{23}R_{13}T_{12} + S_{23}T_{13}U_{12}$$

and similarly for the primed equations with respect to R' . There is an evident mirror-like symmetry between the primed and unprimed equations in the theorem.

In the following examples we describe some special choices of operators satisfying the conditions formulated in Theorem 2.7.

Example 2.8. The following choice of endomorphisms satisfies the assumptions of Theorem 2.7 : $\mathcal{R}$ and $\mathcal{R}'$ are arbitrary Yang-Baxter operators, $\mathcal{U}$ and $\mathcal{S}$ are given by $\mathcal{U}(x \otimes y) = y \otimes x$, $\mathcal{S}(y \otimes x) = x \otimes y$, $x \in X$, $y \in Y$, and $\mathcal{T} = 0$. We denote in this case the resulting operator by $\mathcal{R} \oplus \mathcal{R}'$ and call it the *direct sum* of the Yang-Baxter operators $\mathcal{R}$ and $\mathcal{R}'$.

The equations (I)-(III)' can be verified easily, we have, for example,

$$(\mathbf{1} \otimes \mathcal{R})(\mathcal{U} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{U})(x \otimes x' \otimes y) = y \otimes \mathcal{R}(x \otimes x') = (\mathcal{U} \otimes \mathbf{1})(\mathbf{1} \otimes \mathcal{U})(\mathcal{R} \otimes \mathbf{1})(x \otimes x' \otimes y),$$

which is (I). The remaining equations (IV)-(VI)' are trivial, since $\mathcal{T} = 0$.

Example 2.9. Let $\mathcal{R}$ and $\mathcal{R}'$ be Yang-Baxter operators, $a \in \mathbf{k}$, and $\mathcal{U}, \mathcal{S}$ be invertible operators such that the conditions (I)-(III)' of Theorem 2.7 are satisfied and such that, moreover,

$$(VII) \quad (\mathbf{1} \otimes \mathcal{R}^2) = a \cdot (\mathbf{1} \otimes \mathcal{R}) + (\mathcal{U} \otimes \mathbf{1})(\mathcal{S} \otimes \mathbf{1}) \text{ on } Y \otimes X^{\otimes 2},$$

$$(VII)' \quad (\mathcal{R}'^2 \otimes \mathbf{1}) = a \cdot (\mathcal{R}' \otimes \mathbf{1}) + (\mathbf{1} \otimes \mathcal{U})(\mathbf{1} \otimes \mathcal{S}) \text{ on } X \otimes Y^{\otimes 2}.$$

Then the operators $\mathcal{R}, \mathcal{R}', \mathcal{U}, \mathcal{S}$ and $\mathcal{T} := a \cdot \mathbf{1}$ satisfy all the equations of Theorem 2.7.

This is almost obvious. Notice that the conditions (IV)-(V)' are satisfied trivially, since $(\mathcal{T} \otimes \mathbf{1}) = (\mathbf{1} \otimes \mathcal{T}) = a \cdot \mathbf{1}^{\otimes 3}$. For the same reason (VII) implies (VI) and (VII)' implies (VI)'.

Example 2.10. Let $\mathcal{R}$ be the Yang-Baxter operator on $\mathbf{k}$ given by $\mathcal{R}(1 \otimes 1) = q$, $\mathcal{R}' = \mathcal{R}$, and $\mathcal{U}$, $\mathcal{S}$ and $\mathcal{T}$ be defined by $\mathcal{U}(1 \otimes 1) = qp^{-1}$, $\mathcal{S} = \mathbf{1}$ and $\mathcal{T}(1 \otimes 1) = q - p^{-1}$, for some invertible $p, q \in \mathbf{k}$. Then it is easy to verify the assumptions formulated in Example 2.9 (with $a = q - p^{-1}$) and our construction gives a two-parameter solution of the QYBE represented by the matrix

$$\begin{pmatrix} q & 0 & 0 & 0 \\ 0 & qp^{-1} & q - p^{-1} & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & q \end{pmatrix}$$

Example 2.11. Let $\mathcal{R}$ and $\mathcal{R}'$ be Yang-Baxter q -Hecke operators, $a = (q - q^{-1})$, $\mathcal{U}$ be an invertible operator satisfying (I) and (I)', and $\mathcal{S} = \mathcal{U}^{-1}$. Then again this choice satisfies the assumptions formulated in Example 2.9. This is obvious: under the assumption $\mathcal{S} = \mathcal{U}^{-1}$, (I) implies (II) & (III) and (I)' implies (II)' & (III)', while (VII) (resp. (VII')) becomes the Hecke-type equation for $\mathcal{R}$ (resp. $\mathcal{R}'$).

Example 2.12. Let $\mathcal{R} = \mathcal{R}'$ be defined by $\mathcal{R}(1 \otimes 1) = q$ and let $\mathcal{U}$, $\mathcal{S}$ be given by $\mathcal{U}(1 \otimes 1) = s \cdot (1 \otimes 1)$, $\mathcal{S}(1 \otimes 1) = s^{-1} \cdot (1 \otimes 1)$, for some invertible $q, s \in \mathbf{k}$. These operators satisfy the conditions of Example 2.11 and our construction gives another two-parameter solution represented by the matrix

$$\begin{pmatrix} q & 0 & 0 & 0 \\ 0 & s & q - q^{-1} & 0 \\ 0 & 0 & s^{-1} & 0 \\ 0 & 0 & 0 & q \end{pmatrix}$$

Another choice of the same type is the following. Let $(X, \mathcal{R}), (Y, \mathcal{R}') \in \mathcal{H}_q$, let $\mathcal{U}$ be given by $\mathcal{U}(x \otimes y) = y \otimes x$, let $\mathcal{S} = \mathcal{U}^{-1}$ and let $\mathcal{T} = (q - q^{-1}) \cdot \mathbf{1}$. This choice gives the glueing $\mathcal{R} \oplus_q \mathcal{R}'$ of Theorem 2.4.

We end this section with a categorical characterization of some constructions above. Discuss first the ‘direct sum’ $\mathcal{R} \oplus \mathcal{R}'$ of Example 2.8. For $(X, \mathcal{R}), (Y, \mathcal{R}') \in \mathcal{YB}$, the natural inclusions $\iota_1 : X \hookrightarrow X \oplus Y$ and $\iota_2 : Y \hookrightarrow X \oplus Y$ induce morphisms (denoted by the same symbols) $\iota_1 : (X, \mathcal{R}) \rightarrow (X \oplus Y, \mathcal{R} \oplus \mathcal{R}')$ and $\iota_2 : (Y, \mathcal{R}') \rightarrow (X \oplus Y, \mathcal{R} \oplus \mathcal{R}')$ in the category $\mathcal{YB}$.

Proposition 2.13. *Let $(X, \mathcal{R})$, $(Y, \mathcal{R}')$ and $(Z, \mathcal{Q})$ be elements of $\mathcal{YB}$ and let $f_1 : (X, \mathcal{R}) \rightarrow (Z, \mathcal{Q})$ and $f_2 : (Y, \mathcal{R}') \rightarrow (Z, \mathcal{Q})$ be morphisms in $\mathcal{YB}$ such that*

$$\mathcal{Q}(f_1(x) \otimes f_2(y)) = f_2(y) \otimes f_1(x), \text{ for all } x \in X \text{ and } y \in Y.$$

Then there exists exactly one map $f : (X \oplus Y, \mathcal{R} \oplus \mathcal{R}') \rightarrow (Z, \mathcal{Q})$ in $\mathcal{YB}$ such that $f_i = f \circ \iota_i$, $i = 1, 2$.

Proof. The map f must necessarily be given by $f(\iota_1(x)) = f_1(x)$ and $f(\iota_2(y)) = f_2(y)$, and if we take it as the definition of f , we easily verify that $\mathcal{S} \circ (f \otimes f) = (f \otimes f)(\mathcal{R} \oplus \mathcal{R}')$. ■

This proposition provides the following categorial characterization of the direct sum $\mathcal{R} \oplus \mathcal{R}'$. Let $\mathcal{D}$ be the category whose objects are quadruples of the form $(V, A, B, \mathcal{Q})$, with $(V, \mathcal{Q}) \in \mathcal{YB}$, A and B vector subspaces of V , $\mathcal{Q}(A \otimes A) \subset A \otimes A$, $\mathcal{Q}(B \otimes B) \subset B \otimes B$, $\mathcal{Q}(a \otimes b) = b \otimes a$ and $\mathcal{Q}(b \otimes a) = a \otimes b$, for $a \in A$ and $b \in B$. The morphisms from $(V, A, B, \mathcal{Q})$ to $(W, A', B', \mathcal{Q}')$ in $\mathcal{D}$ are morphisms $f : (V, \mathcal{Q}) \rightarrow (W, \mathcal{Q}')$ in $\mathcal{YB}$ such that $f(A) \subset A'$ and $f(B) \subset B'$. Define the functor $\Delta : \mathcal{D} \rightarrow \mathcal{YB} \times \mathcal{YB}$ by

$$\Delta(V, A, B, \mathcal{Q}) = (A, \mathcal{Q}) \times (B, \mathcal{Q}).$$

Then we have the following characterization (compare the characterization of direct sums as it is given in [7]).

Proposition 2.14. *The functor $\oplus : \mathcal{YB} \times \mathcal{YB} \rightarrow \mathcal{D}$, $(X, \mathcal{R}) \times (Y, \mathcal{R}') \mapsto (X \oplus Y, X, Y, \mathcal{R} \oplus \mathcal{R}')$, is a left adjoint to the functor $\Delta : \mathcal{D} \rightarrow \mathcal{YB} \times \mathcal{YB}$.*

Proof. The proof is fully routine one, and we add it only because we do not suppose that this kind of arguments is commonly known among the readers of this paper.

We must show that, for any $\xi = (V, A, B, \mathcal{Q}) \in \mathcal{D}$ and $\eta = (X, \mathcal{R}) \times (Y, \mathcal{R}') \in \mathcal{YB} \times \mathcal{YB}$, we have a functorial isomorphism

$$\text{Hom}_{\mathcal{D}}(\oplus \eta, \xi) \cong \text{Hom}_{\mathcal{YB} \times \mathcal{YB}}(\eta, \Delta \xi)$$

or, expanding the expressions,

$$\text{Hom}_{\mathcal{D}}((X \oplus Y, X, Y, \mathcal{R} \oplus \mathcal{R}'), (V, A, B, \mathcal{Q})) \cong \text{Hom}_{\mathcal{YB}}((X, \mathcal{R}), (A, \mathcal{Q})) \times \text{Hom}_{\mathcal{YB}}((Y, \mathcal{R}'), (B, \mathcal{Q})).$$

But this is almost clear. An element from the left-hand side Hom-set is given by a linear map $f : X \oplus Y \rightarrow V$. The maps $f_i := f \circ \iota_i$, $i = 1, 2$, define the morphisms (denoted by the same symbols) $f_1 : (X, \mathcal{R}) \rightarrow (A, \mathcal{Q})$ and $f_2 : (Y, \mathcal{R}') \rightarrow (B, \mathcal{Q})$ from $\mathcal{YB}$, thus $f_1 \times f_2$ is an element of the right-hand side Hom-set.

On the other hand, two such maps f_1 and f_2 give rise to the element f from the left-hand side Hom-set exactly as it was shown in Proposition 2.14. ■

Let us now consider the glueing $\mathcal{R} \oplus_q \mathcal{R}'$ in the same categorical way. We again have the natural inclusions $j_1 : X \rightarrow X \oplus Y$ and $j_2 : Y \rightarrow X \oplus Y$ which induce, by $\iota_1(x \otimes x') := j_1(x) \otimes j_1(x')$ and $\iota_2(y \otimes y') := j_2(y) \otimes j_2(y')$ the morphisms (denoted by the same symbols) $\iota_1 : (X, \mathcal{R}) \rightarrow (X \oplus Y, \mathcal{R} \oplus_q \mathcal{R}')$ and $\iota_2 : (Y, \mathcal{R}') \rightarrow (X \oplus Y, \mathcal{R} \oplus_q \mathcal{R}')$ in $\mathcal{H}_q$. We then have the following proposition.

Proposition 2.15. *Let $(X, \mathcal{R})$, $(Y, \mathcal{R}')$ and $(Z, \mathcal{Q})$ be elements of $\mathcal{H}_q$, let $f_1 : (X, \mathcal{R}) \rightarrow (Z, \mathcal{Q})$ and $f_2 : (Y, \mathcal{R}') \rightarrow (Z, \mathcal{Q})$ be morphisms in $\mathcal{H}_q$, and let $\phi : X \otimes Y \rightarrow Z$ be a linear map. Then there exists exactly one morphism $f : (X \oplus Y, \mathcal{R} \oplus_q \mathcal{R}') \rightarrow (Z, \mathcal{Q})$ in $\mathcal{H}_q$ such that*

$$f \circ \iota_1 = f_1, \quad f \circ \iota_2 = f_2, \quad f(j_1(x) \otimes j_2(y)) = \phi(x \otimes y),$$

for any $x \in X$ and $y \in Y$.

Proof. We identify, for simplicity, the elements of X (resp. Y) with their images under j_1 (resp. j_2) in $X \oplus Y$. We already know that

$$f|_{X \otimes X} = f_1, \quad f|_{Y \otimes Y} = f_2, \quad f|_{X \otimes Y} = \phi.$$

Suppose that $f|_{Y \otimes X} = \psi$ for some linear map $\psi : Y \otimes X \rightarrow Z$. The basic equation

$$\mathcal{Q} \circ f = f \circ (\mathcal{R} \oplus_q \mathcal{R}')$$

is evidently satisfied on $X \otimes X$ and $Y \otimes Y$. For $x \in X$ and $y \in Y$ it gives

$$\mathcal{Q} \circ \phi(x, y) = \psi(y, x), \quad \mathcal{Q} \circ \psi(y, x) = \phi(x, y) + (q - q^{-1}) \cdot \psi(y, x).$$

We can take the first equation above as the definition of ψ , i.e. put $\psi = \mathcal{Q} \circ \phi$. The second equation then reads

$$\mathcal{Q}^2 \circ \phi = \phi - (q - q^{-1}) \cdot \mathcal{Q} \circ \phi$$

and it is clearly a consequence of the Hecke equation for $\mathcal{Q}$. ■

Again we can convert the previous example into a categorial characterization of the glueing. Let $\mathcal{E}$ be the category whose objects are couples $((X, \mathcal{R}), (Y, \mathcal{R}'))$ of elements of $\mathcal{H}_q$ and the

morphisms between $((X, \mathcal{R}), (Y, \mathcal{R}'))$ and $((X_1, \mathcal{R}_1), (Y_1, \mathcal{R}'_1))$ in $\mathcal{E}$ are triples (f, g, ϕ) , with $f \in \text{Hom}_{\mathcal{H}_q}((X, \mathcal{R}), (X_1, \mathcal{R}_1))$, $g \in \text{Hom}_{\mathcal{H}_q}((Y, \mathcal{R}'), (Y_1, \mathcal{R}'_1))$, and $\phi : X \otimes Y \rightarrow (X_1 \oplus Y_1)^{\otimes 2}$ an arbitrary linear map. Define the functor $\Delta : \mathcal{H}_q \rightarrow \mathcal{E}$ by $\Delta((X, \mathcal{R})) = ((X, \mathcal{R}), (X, \mathcal{R}))$. The proof of the following proposition is almost identical with the proof of Proposition 2.15 and we omit it.

Proposition 2.16. *The functor $\oplus_q : \mathcal{E} \rightarrow \mathcal{H}_q$ sending $((X, \mathcal{R}), (Y, \mathcal{R}')) \rightarrow (X \oplus Y, \mathcal{R} \oplus_q \mathcal{R}')$, is a left adjoint to the functor $\Delta : \mathcal{H}_q \rightarrow \mathcal{E}$ defined above.*

3. Glueing of Quantum Matrices.

We recall that a bialgebra over $\mathbf{k}$ is (A, Δ, ϵ) where A is a unital algebra and $\Delta : A \rightarrow A \otimes A$ a co-associative algebra homomorphism forming a coalgebra with counit $\epsilon : A \rightarrow \mathbf{k}$.

Associated to every matrix $R \in M_n \otimes M_n$ is a bialgebra $A(R)$ defined as follows. We take a n^2 generators $\mathbf{t} = (t^i_j)$ regarded as a matrix, and relations [3] cf [2]

$$(6) \quad R(\mathbf{t} \otimes \mathbf{1})(\mathbf{1} \otimes \mathbf{t}) = (\mathbf{1} \otimes \mathbf{t})(\mathbf{t} \otimes \mathbf{1})R, \quad \text{i.e.,} \quad R\mathbf{t}_1\mathbf{t}_2 = \mathbf{t}_2\mathbf{t}_1R$$

in a standard compact notation. The coproduct and counit are

$$(7) \quad \Delta t^i_j = t^i_a \otimes t^a_j, \quad \epsilon t^i_j = \delta^i_j, \quad \text{i.e.,} \quad \Delta \mathbf{t} = \mathbf{t} \otimes \mathbf{t}, \quad \epsilon \mathbf{t} = \mathbf{1}.$$

This structure is well-known and is commonly called a *quantum matrix algebra*. One assumes that R is the matrix representing an invertible Yang-Baxter operator in order to be sure that the quantum matrix algebra has canonical matrix representations and other nice properties. In this case it is quasi-commutative in the sense of a dual-quasitriangular structure induced by R cf [18].

Further quotienting of these $A(R)$ for standard solutions R gives the matrix description of the quantum function algebras $\mathcal{O}_q(G)$ deforming the ring of representative functions on the standard semisimple Lie groups [3]. The standard $\mathcal{R}_{su(n)}$ solutions give the quantum matrices $M_q(n)$ and the quotient of these by a q -determinant relation gives the matrix quantum group $SL_q(n)$ or (in a suitable context) $SU_q(n)$. One can also localize the q -determinant and obtain the matrix quantum group $GL_q(n)$. In all cases these algebras are like the ring of functions on some ‘quantum group’ from the point of view of non-commutative geometry.

On the other hand, the $A(R)$ bialgebra construction is not limited to these standard Yang-Baxter operators and in general need not have any such determinants etc. Thus for the general

theory we are limited to the quantum matrix setting. We give in this section a description of $A(R \oplus_q R')$ where $R \oplus_q R'$ as from (3). In order to do this we will be led to introduce the notion of a rectangular quantum matrix algebra. These include the quantum matrix algebras above as well as the quantum spaces of the next section.

Definition 3.1. Let $R \in M_m \otimes M_m$ and $R' \in M_n \otimes M_n$ be the matrices. We define the $m \times n$ rectangular quantum matrix algebra $A(R : R')$ to be the algebra generated by $\mathbf{1}$ and $\mathbf{t} = (t^i_{j'})$ for $1 \leq i \leq m$ and $1 \leq j' \leq n$ with relations

$$R(\mathbf{t} \otimes \mathbf{1})(\mathbf{1} \otimes \mathbf{t}) = (\mathbf{1} \otimes \mathbf{t})(\mathbf{t} \otimes \mathbf{1})R', \quad \text{i.e.,} \quad R\mathbf{t}_1\mathbf{t}_2 = \mathbf{t}_2\mathbf{t}_1R'$$

We use here the same compact notation as above for quantum matrices. If one wants to see the relations written completely explicitly in terms of the individual generators, they are

$$(8) \quad R^i_a{}^k{}_b t^a_{j'} t^b_{l'} = t^k_{b'} t^i_{a'} R^{a'}_{j'}{}^{b'}_{l'}, \quad \forall 1 \leq i, k \leq m, 1 \leq j', l' \leq n$$

where summation is over $1 \leq a, b \leq m$ and $1 \leq a', b' \leq n$. We use unprimed indices for the range $1, \dots, m$ and primed indices for the range $1, \dots, n$. An elementary proposition is the following.

Proposition 3.2. Let $R \in M_m \otimes M_n$ and $R' \in M_n \otimes M_n$ be matrices and let

$$\mathbf{a} = (a^i_j \in A(R)), \quad \mathbf{b} = (b^i_{j'} \in A(R : R')), \quad \mathbf{c} = (c^{i'}_j \in A(R' : R)), \quad \mathbf{d} = (d^{i'}_{j'} \in A(R'))$$

denote the generators of the corresponding square and rectangular quantum matrices as shown. The indices run in the ranges described in the conventions above. Then there are algebra homomorphisms

1. $A(R) \rightarrow A(R : R') \otimes A(R' : R), \quad a^i_j \mapsto b^i_{a'} \otimes c^{a'}_j$
2. $A(R') \rightarrow A(R' : R) \otimes A(R : R'), \quad d^{i'}_{j'} \mapsto c^{i'}_a \otimes b^a_{j'}$
3. $A(R : R') \rightarrow A(R) \otimes A(R : R'), \quad b^i_{j'} \mapsto a^i_a \otimes b^a_{j'}$
4. $A(R : R') \rightarrow A(R : R') \otimes A(R'), \quad b^i_{j'} \mapsto b^i_{a'} \otimes d^{a'}_{j'}.$

Proof. The proof is easy in the compact notation. Thus $A(R : R') \otimes A(R' : R)$ is generated by $\mathbf{1}$ and $\mathbf{b}, \mathbf{c}$ which mutually commute. We have to show that $\mathbf{a} \mapsto \mathbf{bc}$ is an algebra homomorphism. Thus $R\mathbf{b}_1\mathbf{c}_1\mathbf{b}_2\mathbf{c}_2 = R\mathbf{b}_1\mathbf{b}_2\mathbf{c}_1\mathbf{c}_2 = \mathbf{b}_2\mathbf{b}_1R'\mathbf{c}_1\mathbf{c}_2 = \mathbf{b}_2\mathbf{b}_1\mathbf{c}_2\mathbf{c}_1R = \mathbf{b}_2\mathbf{c}_2\mathbf{b}_1\mathbf{c}_1R$ as required for the first map. The proofs for the other maps are similar. ■

This proposition extends the notion of quantum linear algebra as in [20] to the setting of rectangular matrices. The first map corresponds to the ability to multiply an $m \times n$ matrix with an $n \times m$ matrix to obtain an $m \times m$ matrix. Similarly for the second map to obtain an $n \times n$ matrix. The third map corresponds to the product of an $m \times m$ matrix with an $m \times n$ matrix from the left. Similarly for the fourth map from the right by an $n \times n$ matrix. These maps are therefore the rectangular counterparts of the coproduct Δ which corresponds to usual square matrix multiplication. The most general statement which covers all these maps as special cases is the following: if R, R' are as above and $R'' \in M_p \otimes M_p$ then there is an algebra homomorphism

$$(9) \quad \Delta_{R,R',R''} : A(R : R'') \rightarrow A(R : R') \otimes A(R' : R''), \quad \mathbf{t} \mapsto \mathbf{b} \otimes \mathbf{c}$$

where $\mathbf{t}$ is a quantum $m \times p$ matrix, $\mathbf{b}$ is $m \times n$ and $\mathbf{c}$ is $n \times p$ and the map corresponds to the matrix multiplication of the latter two to obtain the former. Moreover, the family of maps *taken together* are coassociative in the sense

$$(10) \quad (\text{id} \otimes \Delta_{R',R'',R'''}) \circ \Delta_{R,R',R'''} = (\Delta_{R,R',R''} \otimes \text{id}) \circ \Delta_{R,R'',R'''}$$

as a map $A(R : R''') \rightarrow A(R : R') \otimes A(R' : R'') \otimes A(R'' : R''')$. This includes all possible coassociativity conditions arising from associativity of rectangular matrix multiplication. For example, it says that Δ in (7) is coassociative and that $A(R : R')$ is a left-right $A(R) - A(R')$ -bicomodule via the last two maps in the proposition. In summary, rectangular quantum matrices are not individually bialgebras but they all fit together into a weaker ‘co-groupoid’ structure on the entire family. We can now present our main result of this section in terms of such objects.

Theorem 3.3. *Let $R \in M_m \otimes M_m$ and $R' \in M_n \otimes M_n$ be the matrices of finite-dimensional q -Hecke Yang-Baxter operators. Then $A(R \oplus_q R')$ is the algebra generated by $A(R), A(R : R'), A(R' : R)$ and $A(R')$ as subalgebras, modulo relations between them. We write the matrix generator of $A(R \oplus_q R')$ as*

$$\mathbf{t} = \begin{pmatrix} \mathbf{a} & \mathbf{b} \\ \mathbf{c} & \mathbf{d} \end{pmatrix}, \quad \text{where,} \quad \mathbf{a} = (a^i_j), \quad \mathbf{b} = (b^i_{j'}), \quad \mathbf{c} = (c^{i'}_j), \quad \mathbf{d} = (d^{i'}_{j'})$$

where $i, j = 1, \dots, m$ and $i', j' = 1 \dots n$ and $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}$ are the respective matrix generators of the subalgebras. These are as in Proposition 3.2 and have explicitly the relations

$$(I) \quad R\mathbf{a}_1\mathbf{a}_2 = \mathbf{a}_2\mathbf{a}_1R, \quad R\mathbf{b}_1\mathbf{b}_2 = \mathbf{b}_2\mathbf{b}_1R', \quad R'\mathbf{c}_1\mathbf{c}_2 = \mathbf{c}_2\mathbf{c}_1R, \quad R'\mathbf{d}_1\mathbf{d}_2 = \mathbf{d}_2\mathbf{d}_1R'.$$

The relations between them are

$$(II) \quad R\mathbf{a}_1\mathbf{b}_2 = \mathbf{b}_2\mathbf{a}_1, \quad \mathbf{c}_1\mathbf{a}_2 = \mathbf{a}_2\mathbf{c}_1R, \quad R'\mathbf{c}_1\mathbf{d}_2 = \mathbf{d}_2\mathbf{c}_1, \quad \mathbf{d}_1\mathbf{b}_2 = \mathbf{b}_2\mathbf{d}_1R'$$

$$(III) \quad \mathbf{b}_1 \mathbf{c}_2 = \mathbf{c}_2 \mathbf{c}_1, \quad \mathbf{a}_1 \mathbf{d}_2 - \mathbf{d}_2 \mathbf{a}_1 = (q^{-1} - q) P \mathbf{c}_1 \mathbf{b}_2$$

where P is the permutation matrix. Moreover, the matrix coalgebra structure in these terms is

$$\Delta \mathbf{a} = \mathbf{a} \otimes \mathbf{a} + \mathbf{b} \otimes \mathbf{c}, \quad \epsilon \mathbf{a} = \mathbf{1}, \quad \Delta \mathbf{b} = \mathbf{b} \otimes \mathbf{d} + \mathbf{a} \otimes \mathbf{c}, \quad \epsilon \mathbf{b} = 0$$

$$\Delta \mathbf{c} = \mathbf{c} \otimes \mathbf{a} + \mathbf{d} \otimes \mathbf{c}, \quad \epsilon \mathbf{c} = 0, \quad \Delta \mathbf{d} = \mathbf{d} \otimes \mathbf{d} + \mathbf{c} \otimes \mathbf{b}, \quad \epsilon \mathbf{d} = \mathbf{1}$$

Proof. We compute the structure of $A(R \oplus_q R')$ from (6) which in explicit component form reads

$$(R \oplus_q R')^I{}_A{}^K{}_B t^A{}_J t^B{}_L = t^K{}_B t^I{}_A (R \oplus_q R')^A{}_J{}^B{}_L, \quad \forall 1 \leq I, J, K, L \leq m+n$$

where the summed indices also run from $1, \dots, m+n$. As in (3) we break the range into $1, \dots, m$ which we represent with lower case labels and the rest which we represent by primed labels running $1, \dots, n$. With these conventions the relations are of 16 types according as the four free labels are each in their upper or lower ranges. In evaluating these we must sum over both types of label so each equation has four possible terms on each side. Note now that the structure of $R \oplus_q R'$ in (3) is such that there are only five types of non-zero entries,

$$(R \oplus_q R')^i{}_j{}^k{}_l = R^i{}_j{}^k{}_l, \quad (R \oplus_q R')^{i'}{}_{j'}{}^{k'}{}_{l'} = R'^{i'}{}_{j'}{}^{k'}{}_{l'}$$

$$(R \oplus_q R')^i{}_j{}^{k'}{}_{l'} = \delta^i{}_j \delta^{k'}{}_{l'}, \quad (R \oplus_q R')^{i'}{}_{j'}{}^k{}_l = \delta^{i'}{}_{j'} \delta^k{}_l, \quad (R \oplus_q R')^i{}_j{}^k{}_l = (q - q^{-1}) \delta^i{}_l \delta^{k'}{}_{j'}.$$

We also write the entries of $t^I{}_J$ in the block form stated and write $Q = R \oplus_q R$ for brevity. Then we evaluate our sixteen relations as

$$(ikjl) : \quad Q^i{}_a{}^k{}_b t^a{}_j t^b{}_l = t^k{}_b t^i{}_a Q^a{}_j{}^b{}_l \quad \text{i.e., } R \mathbf{a}_1 \mathbf{a}_2 = \mathbf{a}_2 \mathbf{a}_1 R$$

$$(ikj'l') : \quad Q^i{}_a{}^k{}_b t^a{}_j t^b{}_{l'} = t^k{}_{b'} t^i{}_a Q^a{}_j{}^{b'}{}_{l'} \quad \text{i.e., } R \mathbf{a}_1 \mathbf{b}_2 = \mathbf{b}_2 \mathbf{a}_1$$

$$(ikj'l) : \quad Q^i{}_a{}^k{}_b t^a{}_j t^b{}_l = t^k{}_b t^i{}_{a'} Q^{a'}{}_{j'}{}^b{}_l + t^k{}_{b'} t^i{}_a Q^a{}_j{}^{b'}{}_{l'} \quad \text{i.e., } R \mathbf{b}_1 \mathbf{a}_2 = \mathbf{a}_2 \mathbf{a}_1 + \mathbf{b}_2 \mathbf{a}_1 (q - q^{-1}) P$$

$$(ikj'l') : \quad Q^i{}_a{}^k{}_b t^a{}_j t^b{}_{l'} = t^k{}_{b'} t^i{}_{a'} Q^{a'}{}_{j'}{}^{b'}{}_{l'} \quad \text{i.e., } R \mathbf{b}_1 \mathbf{b}_2 = \mathbf{b}_2 \mathbf{b}_1 R'$$

$$(ik'jl) : \quad Q^i{}_a{}^{k'}{}_{b'} t^a{}_j t^{b'}{}_{l'} + Q^i{}_{a'}{}^{k'}{}_b t^{a'}{}_{j'} t^b{}_l = t^{k'}{}_{b'} t^i{}_a Q^a{}_j{}^b{}_l \quad \text{i.e., } \mathbf{a}_1 \mathbf{c}_2 + (q - q^{-1}) P \mathbf{c}_1 \mathbf{a}_2 = \mathbf{c}_2 \mathbf{a}_1 R$$

$$(ik'j'l') : \quad Q^i{}_a{}^{k'}{}_{b'} t^a{}_j t^{b'}{}_{l'} + Q^i{}_{a'}{}^{k'}{}_b t^{a'}{}_{j'} t^b{}_{l'} = t^{k'}{}_{b'} t^i{}_{a'} Q^{a'}{}_{j'}{}^b{}_{l'} \quad \text{i.e., } \mathbf{a}_1 \mathbf{d}_2 + (q - q^{-1}) P \mathbf{c}_1 \mathbf{b}_2 = \mathbf{d}_2 \mathbf{a}_1$$

$$(ik'j'l) : \quad Q^i{}_a{}^{k'}{}_{b'} t^a{}_j t^{b'}{}_l + Q^i{}_{a'}{}^{k'}{}_b t^{a'}{}_{j'} t^b{}_l = t^{k'}{}_{b'} t^i{}_a Q^a{}_j{}^{b'}{}_l + t^{k'}{}_b t^i{}_{a'} Q^{a'}{}_{j'}{}^b{}_l$$

$$\text{i.e., } \mathbf{b}_1 \mathbf{c}_2 + (q - q^{-1}) P \mathbf{d}_1 \mathbf{a}_2 = (q - q^{-1}) \mathbf{d}_2 \mathbf{a}_1 P + \mathbf{c}_2 \mathbf{b}_1$$

$$(ik'j'l') : \quad Q^i{}_a{}^{k'}{}_{b'} t^a{}_j t^{b'}{}_{l'} + Q^i{}_{a'}{}^{k'}{}_b t^{a'}{}_{j'} t^{b'}{}_{l'} = t^{k'}{}_{b'} t^i{}_{a'} Q^{a'}{}_{j'}{}^{b'}{}_{l'}$$

$$\text{i.e., } \mathbf{b}_1 \mathbf{d}_2 + (q - q^{-1}) P \mathbf{d}_1 \mathbf{b}_2 = \mathbf{d}_2 \mathbf{b}_1 R'$$

and eight similar relations. The $(ikjl), (ikjl'), (ikj'l'), (ik'jl')$ relations here are as stated while the $(ikj'l)$ relation is equivalent to the $(ikjl')$ one because R is q -Hecke. In matrix terms this means $R_{21}R = 1 + (q - q^{-1})PR$ where P is the permutation matrix. Remembering that our suffices in the compact notation relate to the position in a matrix tensor product, the action by conjugation by P is to permute the relevant suffices. Thus $PRP = R_{21}$ etc. as needed in this computation. The action of P also means that the $(ik'j'l)$ also simplifies to $\mathbf{b}_1\mathbf{c}_2 = \mathbf{c}_2\mathbf{b}_1$ while the Hecke property for R and R' means in the same way as above that $(ik'jl)$ and $(ik'j'l')$ are equivalent to $\mathbf{c}_1\mathbf{a}_2 = \mathbf{a}_2\mathbf{c}_1R$ and $\mathbf{d}_1\mathbf{b}_2 = \mathbf{b}_2\mathbf{d}_1R'$ respectively. The remaining eight relations are reduced in the same way and give the three remaining relations in (I)-(II). The form of the coproduct is from $\Delta t^I_J = t^I_A \otimes t^A_J = t^I_a \otimes t^a_J + t^I_{a'} \otimes t^{a'}_J$ giving four equations according to the range of I, J . The counit is immediate. $\blacksquare$

We can think of this theorem as defining a glueing construction $A(R) *_{q} A(R')$ for quantum matrices consisting of viewing them on the diagonal of a 2×2 block and adjoining the relevant off-diagonal rectangular matrices. In the present case the 2×2 blocks obey a blocked version of the usual $M_q(2)$ algebra. Thus,

Example 3.4. Let $R = (q) = R'$ the one-dimensional q -Hecke Yang-Baxter operator. Then a, b, c, d are 1×1 matrices of generators, relations (I) are empty and (II)-(III) reduce to the standard commutation relations

$$qab = ba, \quad ca = acq, \quad qcd = dc, \quad db = dbq, \quad bc = cb, \quad ad - da = (q^{-1} - q)cb$$

for the 2×2 quantum matrices $M_q(2)$. This fits with Example 2.5 in Section 2.

Example 3.5. Let $R = (q)$ and $R' = (-q^{-1})$ as in Example 2.6 and $q^2 \neq -1$. Then a, b, c, d are 1×1 matrices and obey the relations

$$b^2 = 0, \quad c^2 = 0$$

$$qab = ba, \quad ca = acq, \quad cd = -qdc, \quad -qdb = bd, \quad bc = cb, \quad ad - da = (q^{-1} - q)cb$$

for the non-standard 2×2 quantum matrices $M_q(1|1)$ as computed in [5]. The first two come from the middle two of (I) as $qbb = bb(-q^{-1})$ and $qcc = cc(-q^{-1})$ and the remainder from (II)-(III).

It is easy to see from the structure in Theorem 3.3 that setting $\mathbf{b} = \mathbf{c} = 0$ gives a bialgebra surjection

$$(11) \quad A(R \oplus_q R') \rightarrow A(R) \otimes A(R').$$

Thus, from (III) we see that $\mathbf{a}$ and $\mathbf{d}$ commute in the quotient as they should in $A(R) \otimes A(R')$ while the other relations either become empty or those of $A(R), A(R')$. This map corresponds to the usual block-diagonal embedding of $M_m \times M_n \subset M_{n+m}$ as semigroups. Thus provided we use our $\oplus_q$ construction when blocking quantum matrices into bigger ones, we have a coherent quantum linear algebra.

For a full understanding of this quantum linear algebra one must understand not only multiplication but addition, to which we turn in the next section at least for the $1 \times n$ or $m \times 1$ cases.

4. Glueing of Quantum-Braided Vector Spaces.

We have introduced rectangular quantum matrix algebras $A(R : R')$ in the last section. When $R' = R$ we have the usual square quantum matrices $A(R)$. On the other hand $A(R : \lambda)$ with $\lambda \in \mathbf{k}$ is the algebra with generators and relations

$$(12) \quad R\mathbf{v}_1\mathbf{v}_2 = \mathbf{v}_2\mathbf{v}_1\lambda, \quad \mathbf{v} = (v^i).$$

This is the algebra of (left-covariant) quantum vectors and denoted $V_L(R)$. Likewise it is clear that $A(\lambda : R)$ is the algebra with generators and relations

$$(13) \quad \lambda\mathbf{x}_1\mathbf{x}_2 = \mathbf{x}_2\mathbf{x}_1R, \quad \mathbf{x} = (x_i).$$

This is the algebra of (right-covariant) quantum covectors and denoted $V^{\sim}(R)$.

The usual quantum planes $\mathbb{C}_q^n$ are of this type with R the matrix representing the operator $\mathcal{R}_{su(n)}$ in Example 2.5. The constructions work best in the case where R are q -Hecke Yang-Baxter operators for in this case one has a braided-coproduct, counit and braiding [15]

$$(14) \quad V^{\sim}(R) = A(R : q), \quad \underline{\Delta}x_i = x_i \otimes 1 + 1 \otimes x_i, \quad \underline{\epsilon}x_i = 0, \quad \Psi(\mathbf{x}_1 \otimes \mathbf{x}_2) = q \mathbf{x}_2 \otimes \mathbf{x}_1 R$$

There is also a braided-antipode $\underline{S}x_i = -x_i$. The notion of braided-Hopf algebras was introduced by one of the authors and we refer to the review [13] for details. The main difference from the usual axioms is that $\underline{\Delta} : V^{\sim}(R) \rightarrow V^{\sim}(R) \underline{\otimes} V^{\sim}(R)$ where $\underline{\otimes}$ is not the usual commuting tensor

product but the braided tensor product algebra

$$(15) \quad (a \otimes b)(c \otimes d) = a\Psi(b \otimes c)d, \quad \forall a, b, c, d \in V^\sim(R).$$

Here Ψ and the other maps extend to products of the generators in a natural way. For a braided-categorical setting (which we defer to the next section) one needs R here to be invertible so that Ψ is invertible. On the other hand this is not essential in our direct algebraic application here.

By allowing ourselves these slightly more general braided-coproduct structures we see that we can express via $\underline{\Delta}$ the addition of two covectors to obtain a new covector, provided we remember the braiding Ψ . This is an important construction with many corollaries. For example, one obtains at once differential calculi on the quantum planes by making infinitesimal additions [17].

In this section we compute $V^\sim(R \oplus_q R)$. There is a strictly analogous theory for vectors also. Let us note first that all our algebras have quadratic relations and hence are naturally $\mathbb{Z}$ -graded by the degree of the generators. Now just as there is a standard $\mathbb{Z}_2$ -graded tensor product of super-algebras, there is a natural braided tensor product of $\mathbb{Z}$ -graded algebras.

Definition 4.1. *cf [14] Let A, B be $\mathbb{Z}$ -graded algebras over $\mathbf{k}$ and let $q \in \mathbf{k}$. We define the associative $\mathbb{Z}$ -graded braided tensor product algebra $A \otimes_q B$ as $A \otimes B$ with product*

$$(a \otimes b)(c \otimes d) = a\Psi_{\mathbb{Z}}(b \otimes c)d, \quad \Psi_{\mathbb{Z}}(b \otimes c) = q^{|b||c|}c \otimes b, \quad \forall a, c \in A, \quad b, d \in B$$

on homogeneous elements of degree $| \cdot |$.

The underlying braided tensor product construction here and in (15) has been introduced by one of the authors as part of the theory of braided groups. See [13] for details. In the present case it is trivial to verify associativity directly.

Proposition 4.2. *Let $R \in M_m \otimes M_m$ and $R' \in M_n \otimes M_n$ be the matrices of q -Hecke Yang-Baxter operators. Then $V^\sim(R \oplus_q R') = V^\sim(R) \otimes_q V^\sim(R')$, the $\mathbb{Z}$ -graded braided tensor product of $V^\sim(R)$ and $V^\sim(R')$.*

Explicitly, we write the covector generator of $V^\sim(R \oplus_q R')$ as $(\mathbf{x}, \mathbf{y})$ where $\mathbf{x} = (x_i)$ and $\mathbf{y} = (y_{i'})$ are the respective covector generators of the subalgebras and have the relations

$$(I) \quad q\mathbf{x}_1\mathbf{x}_2 = \mathbf{x}_2\mathbf{x}_1R, \quad q\mathbf{y}_1\mathbf{y}_2 = \mathbf{y}_2\mathbf{y}_1R'$$

as above. The $\mathbb{Z}$ -graded braided tensor product is equivalent to the additional cross relations

$$(II) \quad \mathbf{y}_1\mathbf{x}_2 = q\mathbf{x}_2\mathbf{y}_1.$$

Moreover, the braided-coproduct restricts to those of $V^\sim(R), V^\sim(R')$ which appear as sub-braided-Hopf algebras.

Proof. The Definition 4.1 of the braided-tensor product for $\mathbb{Z}$ -graded algebras immediately gives the commutation relations (II) between the factors. Conversely, these commutation relations can be used to put all the $\mathbf{x}$ generators to the left of the $\mathbf{y}$ generators, from which it follows that this description is equivalent to the form of $\otimes_q$ in Definition 4.1. To compute $V^\sim(R \oplus_q R')$ directly we use the same conventions for the range of indices as in the proof of Theorem 3.3, so our relations are explicitly

$$qx_Jx_L = x_Bx_A(R \oplus_q R')^A{}_J{}^B{}_L, \quad \forall 1 \leq J, L \leq m+n.$$

We have four types of equation according to the range of (JL) and in each one we have up to four terms on each side according to the breakdown of the range of (AB) . Remembering the explicit form of $Q = R \oplus_q R'$ from Theorem 3.3 and denoting the lower and upper ranges by $\{x_i\} = \mathbf{x}$ and $\{x_{j'}\} = \mathbf{y}$, we have

$$(jl) : \quad qx_Jx_L = x_Bx_AQ^a{}_j{}^b{}_l, \quad \text{i.e., } q\mathbf{x}_1\mathbf{x}_2 = \mathbf{x}_2\mathbf{x}_1R$$

$$(jl') : \quad qx_Jx_{l'} = x_{b'}x_aQ^a{}_j{}^{b'}{}_{l'}, \quad \text{i.e., } q\mathbf{x}_1\mathbf{y}_2 = \mathbf{y}_2\mathbf{x}_1$$

$$(j'l) : \quad qx_{j'}x_l = x_bx_{a'}Q^{a'}{}_{j'}{}^b{}_l + x_{b'}x_aQ^a{}_j{}^{b'}{}_{l'}, \quad \text{i.e., } q\mathbf{y}_1\mathbf{x}_2 = \mathbf{x}_2\mathbf{y}_1 + \mathbf{y}_2\mathbf{x}_1(q - q^{-1})P$$

$$(j'l') : \quad qx_{j'}x_{l'} = x_{b'}x_{a'}Q^{a'}{}_{j'}{}^{b'}{}_{l'}, \quad \text{i.e., } q\mathbf{y}_1\mathbf{y}_2 = \mathbf{y}_2\mathbf{y}_1R'$$

as required. The third equation here is redundant.

The braided-coproduct of $V^\sim(R \oplus_q R')$ is by definition the linear one as in (14) on the generators but (as explained) it extended to products of the generators as an algebra homomorphism to the braided tensor product $V^\sim(R \oplus_q R') \underline{\otimes} V^\sim(R \oplus_q R')$. The latter is defined as in (15) with the operator Ψ induced by the Yang-Baxter operator $\mathcal{R} \oplus_q \mathcal{R}'$. But from its definition in Theorem 2.4 we know that this operator restricts on the generators x_i and $y_{i'}$ to $\mathcal{R}$ and $\mathcal{R}'$ respectively. Likewise for their extensions to products of the generators. Hence the braided-coproduct $\underline{\Delta}$ restricted to $V^\sim(R)$ and $V^\sim(R')$ coincides with their braided-coproducts whose extension to products is defined via $\mathcal{R}, \mathcal{R}'$. ■

This proposition was the original motivation for the $\oplus_q$ construction. Namely, one can ask how to build up higher-dimensional quantum space algebras by tensor product of lower-dimensional ones, corresponding from the point of view of non-commutative geometry to the usual cartesian

product. Unfortunately, in any fixed braided category, there is no canonical notion of tensor product of braided-Hopf algebras or braided-bialgebras. Our answer is that each of our quantum spaces comes with its own braiding Ψ as determined from its associated Yang-Baxter matrix R . At least in the q -Hecke setting one can q -tensor product them, providing their associated braidings are also glued by the $\oplus_q$ construction of Section 2.

Example 4.3. Let R, R' represent the Yang-Baxter operators $\mathcal{R}_{su(m)}$ and $\mathcal{R}_{su(n)}$ respectively, as in Example 2.5. The associated quantum-covector algebras $\mathbb{C}_q^m$ and $\mathbb{C}_q^n$ are given by

$$x_j x_i = q x_i x_j, \quad \forall 1 \leq i < j \leq m; \quad y_{j'} y_{i'} = q y_{i'} y_{j'}, \quad \forall 1 \leq i' < j' \leq n.$$

Their $\otimes_q$ is of the same form with covector generator $(\mathbf{x}, \mathbf{y})$ and similar relations because our ordering of the basis vectors is with all the primed generators $y_{i'}$ appearing as $x_{m+i'}$ in $\mathbb{C}_q^{m+n}$ and hence with labels greater than the unprimed ones. Thus

$$\mathbb{C}_q^{m+n} = \mathbb{C}_q^m \otimes_q \mathbb{C}_q^n$$

as required from Example 2.5 and the above proposition.

Example 4.4. Let $R = (q)$ and $R' = (-q^{-1})$ be the two 1-dimensional q -Hecke Yang-Baxter operators as in Example 2.6. Then $V^{\sim}(R \oplus_q R')$ is the algebra

$$qxx = xqx, \quad qyy = -q^{-1}yy, \quad yx = qxy.$$

The first relation is empty and the second is $y^2 = 0$ if $q^2 \neq -1$. The remaining relation comes from the braided tensor product $\otimes_q$. This is the usual non-standard plane associated to the R -matrix in Example 2.6.

We have described here the so-called ‘bosonic’ covectors which appear in our general setting as the $1 \times n$ matrices $A(q : R)$ and studied how it behaves under $\oplus_q$ in the second input. There are also the so-called ‘fermionic’ covectors $A(-q^{-1} : R)$ in our notation. Their behaviour under $\oplus_q$ is the same as in Proposition 4.2 with q replaced everywhere by $-q^{-1}$. Likewise there are the analogous vector cases $A(R : q)$ and $A(R : -q^{-1})$. All of these important cases can be collected into the following general statement about the behaviour of $A(R : R')$ under $\oplus_q$ in each input.

Proposition 4.5. *Let $R \in M_m \otimes M_m$, $R' \in M_n \otimes M_n$ and $R'' \in M_p \otimes M_p$ be the matrices of q -Hecke Yang-Baxter operators. Then*

$$1. A(R : R' \oplus_q R'') = A(R : R')_{R \underline{\otimes}} A(R : R'')$$

$$2. A(R \oplus_q R' : R'') = A(R : R'')_{\underline{\otimes}_{R''}} A(R' : R'')$$

where $R \underline{\otimes}$ and $\underline{\otimes}_{R''}$ are the braided tensor product algebras defined by the cross relations

$$\mathbf{b} \in A(R : R'), \quad \mathbf{c} \in A(R : R''), \quad R\mathbf{b}_1\mathbf{c}_2 = \mathbf{c}_2\mathbf{b}_1$$

$$\mathbf{b} \in A(R : R''), \quad \mathbf{c} \in A(R' : R''), \quad \mathbf{c}_1\mathbf{b}_2 = \mathbf{b}_2\mathbf{c}_1R''$$

respectively.

Proof. The strategy for $A(R : R' \oplus_q R'')$ is the same as in Proposition 4.2 with the role of q replaced by R . Explicitly decomposing $R' \oplus_q R''$ along the lines in the proof of Theorem 3.3 and putting into the relations

$$R^i_a{}^k{}_b t^a{}_J t^b{}_L = t^k{}_B t^i{}_A (R' \oplus_q R'')^A{}_J{}^B{}_L, \quad \forall 1 \leq i, k \leq m, \quad \forall 1 \leq J, L \leq n + p.$$

for the $m \times (n + p)$ quantum matrices $\mathbf{t} = (\mathbf{b}, \mathbf{c})$, we obtain the four equations

$$R\mathbf{b}_1\mathbf{b}_2 = \mathbf{b}_2\mathbf{b}_1R', \quad R\mathbf{b}_1\mathbf{c}_2 = \mathbf{c}_2\mathbf{b}_1$$

$$R\mathbf{c}_1\mathbf{b}_2 = \mathbf{b}_2\mathbf{c}_1 + \mathbf{c}_2\mathbf{b}_1(q - q^{-1})P, \quad R\mathbf{c}_1\mathbf{c}_2 = \mathbf{c}_2\mathbf{c}_1R''.$$

The third relation here reduces to the second just because R is q -Hecke so that $R_{21}R = 1 + (q - q^{-1})PR$ and since $P\mathbf{c}_2\mathbf{b}_1 = \mathbf{c}_1\mathbf{b}_2P$. Recall that P always denotes the relevant permutation operator in a matrix tensor product. The second relation expresses the non-commutativity in the braided tensor product algebra $R \underline{\otimes}$ defined as in (15) and Definition 4.1 but with braiding

$$\Psi(\mathbf{c}_1 \otimes \mathbf{b}_2) = R_{21}\mathbf{b}_2 \otimes \mathbf{c}_1.$$

The proof for $A(R \oplus_q R' : R'')$ is entirely similar with decomposition $\mathbf{t} = \begin{pmatrix} \mathbf{b} \\ \mathbf{c} \end{pmatrix}$ of an $(m + n) \times p$ quantum matrix. ■

One also has a braided addition law for all quantum matrices $A(R : R')$ when R, R' are the matrices of q -Hecke Yang-Baxter operators,

$$A(R : R') : \quad \underline{\Delta}_{R, R'}\mathbf{t} = \mathbf{t} \otimes 1 + 1 \otimes \mathbf{t}, \quad \underline{\epsilon}\mathbf{t} = 0, \quad \Psi(\mathbf{t}_1 \otimes \mathbf{t}_2) = R_{21}\mathbf{t}_2 \otimes \mathbf{t}_1R'$$

forming a braided-Hopf algebra. This precisely generalizes the addition law (14) from $1 \times n$ to a general $m \times n$ quantum matrix. For example it means that the usual square quantum matrix

algebra $A(R)$ has this linear coproduct or ‘coaddition’ $\underline{\Delta}$ provided one uses the notion of a braided-Hopf algebra. The coaddition is codistributive with respect to the product of quantum matrices in the sense

$$(\text{id} \otimes \cdot) \circ \tau_{23} \circ (\Delta_{R,R',R''} \otimes \Delta_{R,R',R''}) \circ \underline{\Delta}_{R,R''} = (\underline{\Delta}_{R,R'} \otimes \text{id}) \circ \Delta_{R,R',R''}$$

$$(\cdot \otimes \text{id}) \circ \tau_{23} \circ (\Delta_{R,R',R''} \otimes \Delta_{R,R',R''}) \circ \underline{\Delta}_{R,R''} = (\text{id} \otimes \underline{\Delta}_{R,R'}) \circ \Delta_{R,R',R''}$$

where τ_{23} denotes usual transposition in the second and third positions. For example, $A(R)$ becomes a ‘quantum ring’ with both its usual comultiplication and the above coaddition in a distributive way. It is easy to see that for trivial R these notions are the usual notions for the ring of matrices, expressed in terms of the algebra of functions on them.

These remarks about addition of quantum matrices are not directly related to our glueing construction and will be developed elsewhere. The coaddition is however, compatible with Proposition 4.5 in the sense that the factors appear as sub-braided-Hopf algebras. This provides one more reason to be interested in the quantum matrices $A(R : R')$ and their glueing. Clearly, we obtain Proposition 4.2 as a corollary from this point of view. On the other hand the quantum-braided planes are the most well-known of these algebras and of independent interest, and for this reason we emphasized them in the above.

5. Some other properties of the glueing.

We show in this section that our $\oplus_q$ operation is well-behaved with respect to a number of constructions related to duals and link invariants. If $\mathcal{R}_{su(1)}$ has these properties, it follows for example that $\mathcal{R}_{su(n)}$ given by the iterated glueing of $\mathcal{R}_{su(1)}$ (see Example 2.5), has automatically these properties, too. Many other properties of $\mathcal{R}_{su(n)}$ can similarly be understood in this way as arising from the naturality of $\oplus_q$.

Let us introduce first some standard notation and terminology. These notions make sense in any monoidal category, though we shall be interested here only in full subcategories of the category $\mathbf{k}\text{-Vect}$ of finite dimensional vector spaces with its usual tensor product and associativity. We recall that a general monoidal category consists of a category $\mathcal{C}$ and a functor $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ with properties analogous to those for $\mathbf{k}\text{-Vect}$. There is a unit object $\underline{1}$ for the tensor product, and related unity and associativity morphisms which we suppress. For $\mathbf{k}\text{-Vect}$ we have $\underline{1} = \mathbf{k}$.

In a monoidal category, there is a notion of the *left dual* of an object V (see for example [1]). This is an object V^* and morphisms $\text{coev}_V : \underline{1} \rightarrow V \otimes V^*$ (the coevaluation) and $\text{ev}_V : V^* \otimes V \rightarrow \underline{1}$

(the evaluation) satisfying

$$(\text{ev}_V \otimes \text{id})(\text{id} \otimes \text{coev}_V) = \text{id} , \quad (\text{id} \otimes \text{ev}_V)(\text{coev}_V \otimes \text{id}) = \text{id}.$$

We will use the shorthand $V^* \dashv V$ to indicate this situation. Sometimes we say also that the pair (ev, coev) is an *adjunction* between V^* and V . A monoidal category for which every object has a left dual is then called *rigid*.

A typical example is the category $\mathbf{k}\text{-Vect}$ with $V^* = \text{Hom}(V, \mathbf{k})$ and usual evaluation. Notice also that for a basis $(x_1, \dots, x_n)$ of V , there always exist the *dual basis* $(x_1^*, \dots, x_n^*)$ characterized by the property that $\text{coev}(1) = \sum x_i \otimes x_i^*$ and $\text{ev}(x_i^* \otimes x_j) = \delta_{ij}$ (the Kronecker delta).

Suppose we have a morphism $f : V \rightarrow W$ and two adjunctions $V^* \dashv V$ and $W^* \dashv W$. Then there is an adjoint or *dual* morphism $f^* : W^* \rightarrow V^*$ defined by

$$f^* = (\text{ev}_W \otimes \text{id})(\text{id} \otimes f \otimes \text{id})(\text{id} \otimes \text{coev}_V).$$

This dualization via ev, coev can be usefully generalized to the case of a morphism $f : X \otimes V \rightarrow W \otimes Y$ for any objects X, Y . In this case we follow [12] and denote the generalized adjoint morphism by $f^\# : W^* \otimes X \rightarrow Y \otimes V^*$. It is defined in the analogous way as

$$f^\# = (\text{ev}_W \otimes \text{id}^2)(\text{id} \otimes f \otimes \text{id})(\text{id}^2 \otimes \text{coev}_V).$$

Finally, we will need the notion of *braided monoidal category*. This notion was formally introduced in category theory in [8] and also arises in the representation theory of quantum groups. A braided monoidal category is by definition a monoidal category $\mathcal{C}$ equipped with a natural isomorphism $\Psi : - \otimes - \rightarrow - \otimes^{\text{op}} -$, where the latter is the opposite tensor product. This Ψ is called the quasisymmetry or *braiding* and is required to obey some properties analogous to those for the usual transposition of vector spaces, namely

$$\Psi_{V,W \otimes Z} = (\text{id} \otimes \Psi_{V,Z}) \circ (\Psi_{V,W} \otimes \text{id}), \quad \Psi_{V \otimes W,Z} = (\Psi_{V,Z} \otimes \text{id}) \circ (\text{id} \otimes \Psi_{W,Z})$$

for any three objects. Here we continue to suppress the associativity morphisms. Note also that we do not assume that $\Psi_{W,V} \circ \Psi_{V,W} = \text{id}$ for all V, W . Naturality means of course that these isomorphisms are functorial, i.e. all morphisms $f : V \rightarrow X$, $g : W \rightarrow Y$ in $\mathcal{C}$ commute with the braiding in the sense

$$\Psi_{X,W} \circ (f \otimes \text{id}_W) = (\text{id}_W \otimes f) \circ \Psi_{V,W}, \quad \Psi_{V,Y} \circ (\text{id}_V \otimes g) = (g \otimes \text{id}_V) \circ \Psi_{V,W}.$$

A rigid braided monoidal category is a rigid monoidal category which is braided. They have been studied in connection with link invariants in [4].

Now it is well-known that an invertible Yang-Baxter operator $\mathcal{R}$ generates a braided monoidal category. One extends the Yang-Baxter operator from its original vector space V to tensor products of V etc in an obvious way. If V has an adjunction $V^* \dashv V$ and $\mathcal{R}, \mathcal{R}^\#$ are invertible then $\mathcal{R}$ defines in a similar way a rigid braided monoidal category (and leads in fact an Abelian one). This was shown by Lyubashenko in his pioneering work [11][12], and is by now well-known. In fact, we will give for our limited purposes a simplified treatment in which one more invertibility is assumed for convenience. Thus we say that a Yang-Baxter operator is *dualisable* or ‘closed’ if V has an adjunction and $\mathcal{R}, \mathcal{R}^\#$ and $(\mathcal{R}^{-1})^\#$ are invertible. In principle, this last assumption is redundant according to the results effectively contained in [12].

Given a dualisable Yang-Baxter operator $\mathcal{R}$ on V , the first step is to note that it extends to a Yang-Baxter operator $\mathcal{R}^e$ say, on $V \oplus V^*$. The extension is cf [11][12]

$$(16) \quad \mathcal{R}^e|_{V \otimes V} := \mathcal{R}, \quad \mathcal{R}^e|_{V \otimes V^*} := \mathcal{R}^{\#^{-1}}, \quad \mathcal{R}^e|_{V^* \otimes V} := \mathcal{R}^{-1\#}, \quad \mathcal{R}^e|_{V^* \otimes V^*} := \mathcal{R}^*.$$

That this obeys the Yang-Baxter equations follows from applying the adjunctions above to the Yang-Baxter equations for $\mathcal{R}$.

In the following theorem we suppose $X^* \dashv X$ and $Y^* \dashv Y$. Let $(X \oplus Y)^* := X^* \oplus Y^*$ and define $\text{coev} : \mathbf{k} \rightarrow (X \oplus Y) \otimes (X \oplus Y)^*$ by $\text{coev}(1) := \text{coev}_X(1) \oplus \text{coev}_Y(1)$ and let $\text{ev} : (X \oplus Y)^* \otimes (X \oplus Y) \rightarrow \mathbf{k}$ be defined by

$$\text{ev}|_{X^* \otimes X} := \text{ev}_X, \quad \text{ev}|_{Y^* \otimes Y} := \text{ev}_Y, \quad \text{ev}|_{(X^* \otimes Y) \oplus (Y^* \otimes X)} := 0.$$

Then $(X \oplus Y)^* \dashv (X \oplus Y)$ by these maps and we have the following theorem.

Theorem 5.1. *Let $(X, \mathcal{R})$ and $(Y, \mathcal{R}')$ be elements of $\mathcal{H}_q$. Then the Yang-Baxter operator $\mathcal{R} \oplus_q \mathcal{R}'$ is dualisable if and only if both $\mathcal{R}$ and $\mathcal{R}'$ are.*

Proof. The proof is very easy. First, let us agree that the symbols $x, \bar{x}, y$ and $\bar{y}$ will denote arbitrary elements of X, X^*, Y and Y^* respectively and let $\mathcal{Q} = \mathcal{R} \oplus_q \mathcal{R}'$. Then we have the following formulas which can be easily obtained immediately from the definitions:

$$\begin{aligned} \mathcal{Q}^\#(\bar{x} \otimes x) &= \mathcal{R}^\#(\bar{x} \otimes x), \\ \mathcal{Q}^\#(\bar{y} \otimes y) &= \mathcal{R}'^\#(\bar{y} \otimes y) + (q - q^{-1}) \cdot \text{ev}_Y(\bar{y} \otimes y) \cdot \text{coev}_X(1), \\ \mathcal{Q}^\#(\bar{x} \otimes y) &= y \otimes \bar{x}, \quad \mathcal{Q}^\#(\bar{y} \otimes x) = x \otimes \bar{y}, \end{aligned}$$

and

$$\begin{aligned}\mathcal{Q}^{-1\#}(\bar{x} \otimes x) &= \mathcal{R}^{-1\#}(\bar{x} \otimes x) - (q - q^{-1}) \cdot \text{ev}(\bar{x} \otimes x) \cdot \text{coev}_Y(1), \\ \mathcal{Q}^{-1\#}(\bar{y} \otimes y) &= \mathcal{R}'^{-1\#}(\bar{y} \otimes y), \\ \mathcal{Q}^{-1\#}(\bar{x} \otimes y) &= y \otimes \bar{x}, \quad \mathcal{Q}^{-1\#}(\bar{y} \otimes x) = x \otimes \bar{y}.\end{aligned}$$

Now it is immediate to see that the generalized adjoints $\mathcal{Q}^\#$ and $\mathcal{Q}^{-1\#}$ are invertible if and only if $\mathcal{R}^\#, \mathcal{R}'^\#, \mathcal{R}^{-1\#}$ and $\mathcal{R}'^{-1\#}$ are invertible. ■

The equation (16) gives the following explicit formulas for the extension of the glueing:

$$(\mathcal{R} \oplus_q \mathcal{R}')^e|_{(X \oplus Y) \otimes (X \oplus Y)} = (\mathcal{R} \oplus_q \mathcal{R}'), \quad (\mathcal{R} \oplus_q \mathcal{R}')^e|_{(X \oplus Y)^* \otimes (X \oplus Y)^*} = (\mathcal{R} \oplus_q \mathcal{R}')^*,$$

$(\mathcal{R} \oplus_q \mathcal{R}')^e|_{(X \oplus Y)^* \otimes (X \oplus Y)}$ is given by

$$\begin{aligned}(\mathcal{R} \oplus_q \mathcal{R}')^e(\bar{x} \otimes x) &= \mathcal{R}^e(\bar{x} \otimes x) - (q - q^{-1}) \cdot \text{ev}_X(\bar{x} \otimes x) \cdot \text{coev}_Y(1), \\ (\mathcal{R} \oplus_q \mathcal{R}')^e(\bar{y} \otimes y) &= \mathcal{R}'^e(\bar{y} \otimes y), \\ (\mathcal{R} \oplus_q \mathcal{R}')^e(\bar{x} \otimes y) &= y \otimes \bar{x}, \quad (\mathcal{R} \oplus_q \mathcal{R}')^e(\bar{y} \otimes x) = x \otimes \bar{y},\end{aligned}$$

and $(\mathcal{R} \oplus_q \mathcal{R}')^e|_{(X \oplus Y) \otimes (X \oplus Y)^*}$ is given by

$$\begin{aligned}(\mathcal{R} \oplus_q \mathcal{R}')^e(x \otimes \bar{x}) &= \mathcal{R}^e(x \otimes \bar{x}), \\ (\mathcal{R} \oplus_q \mathcal{R}')^e(y \otimes \bar{y}) &= \mathcal{R}'^e(y \otimes \bar{y}) - (q - q^{-1}) \cdot \text{ev}_Y(\mathcal{R}'^e(y \otimes \bar{y})) \cdot \mathcal{R}^e(\text{coev}_X(1)), \\ (\mathcal{R} \oplus_q \mathcal{R}')^e(x \otimes \bar{y}) &= \bar{y} \otimes x, \quad (\mathcal{R} \oplus_q \mathcal{R}')^e(y \otimes \bar{x}) = \bar{x} \otimes y.\end{aligned}$$

Example 5.2. The Yang-Baxter operator $\mathcal{R}_{su(1)}$ on $\mathbf{k} \otimes \mathbf{k}$ given by $\mathcal{R}_{su(1)}(1 \otimes 1) := q \cdot (1 \otimes 1)$, $q \neq 0$, is dualisable, we have clearly

$$\mathcal{R}_{su(1)}^\# = \text{multiplication by } q, \quad \mathcal{R}_{su(1)}^{-1\#} = \text{multiplication by } q^{-1}.$$

As we saw in Example 2.5, the classical Yang-Baxter operator $\mathcal{R}_{su(n)}$ is, for $n \geq 1$, build inductively from $\mathcal{R}_{su(1)}$ and Theorem 5.1 says that this operator is dualisable. It follows from the same kind of arguments that also the non-standard Yang-Baxter operator $\mathcal{R}_{gl(1|1)}$ from Example 2.6 is dualisable.

Now we come to more categorical considerations. Thus suppose that we are given a Yang-Baxter operator $\mathcal{R}$ on V such that it extends to a rigid braided monoidal category containing V , with

$\Psi_{V,V} = \mathcal{R}$. It is immediate that $\mathcal{R}$ is dualisable and $\Psi_{V,V^*}, \Psi_{V^*,V^*}, \Psi_{V^*,V}$ are provided by $\mathcal{R}^e$ above. Some other immediate observations can be obtained at once in the form of the following elementary propositions.

Proposition 5.3. *Suppose that $\mathcal{R}$ extends to the structure of a braided rigid category with $\Psi_{V,V} = \mathcal{R}$ and let $V^* \dashv V$ and $V^{**} \dashv V^*$ be the adjunctions with their associated evaluation and coevaluation maps. Put*

$$\begin{aligned} u_V &:= (\text{ev}_V \otimes \text{id})(\Psi_{V,V^*} \otimes \text{id})(\text{id} \otimes \text{coev}_{V^*}) : V \rightarrow V^{**} \\ v_V &:= (\text{ev}_{V^*} \otimes \text{id})(\text{id} \otimes \Psi_{V,V^*})(\text{id} \otimes \text{coev}_V) : V^{**} \rightarrow V. \end{aligned}$$

Then u_V and v_V are invertible morphisms and their inverses are given by

$$\begin{aligned} u_V^{-1} &= (\text{id} \otimes \text{ev}_{V^*})(\Psi_{V^{**},V} \otimes \text{id})(\text{id} \otimes \text{coev}_V) \\ v_V^{-1} &= (\text{ev}_V \otimes \text{id})(\text{id} \otimes \Psi_{V^{**},V})(\text{coev}_{V^*} \otimes \text{id}). \end{aligned}$$

Proposition 5.4. *Let $\mathcal{R}$ extend to the structure of a rigid braided monoidal category as in Proposition 5.3, and define by induction $V^{[0]} := V$ and $V^{[i+1]} := (V^{[i]})^*$ for $i \geq 0$. The braiding has the following properties.*

1. $\Psi_{V,V}$ is a dualisable Yang-Baxter operator,
2. $\Psi_{V^{[i+1]},V^{[j+1]}} = \left(\Psi_{V^{[i]},V^{[j]}}\right)^*$, $i, j \geq 0$,
3. $\Psi_{V^{[i+2]},V} = (\text{id} \otimes v_{V^{[i]}}^{-1})\Psi_{V^{[i]},V}(v_{V^{[i]}} \otimes \text{id})$, for $i \geq 0$,
4. $\Psi_{V,V^{[i+2]}} = (v_{V^{[i]}}^{-1} \otimes \text{id})\Psi_{V,V^{[i]}}(\text{id} \otimes v_{V^{[i]}})$, for $i \geq 0$.

The last two equations hold also with u_V in place of v_V .

Notice that equations 3. and 4. of this proposition say exactly that the maps $v_{V^{[i]}}$ commute with the braiding, as they must since they are morphisms. That they are morphisms is evident since, by assumption, ev, coev are morphisms. Notice also that Proposition 5.4 together with formula (16) enables one to express inductively $\Psi_{V^{[i]},V^{[j]}}$, for any $i, j \geq 0$, via $\Psi_{V,V}$. Thus we arrive at the following proposition.

Proposition 5.5. *(cf [11][12]) Let V be a vector space with adjunction V^* and denote by $\mathcal{C}(V, V^*)$ the full rigid subcategory of $\mathbf{k}\text{-Vect}$ generated by V, V^* . Let $(V, \mathcal{R})$ be a Yang-Baxter operator. Then $\mathcal{R}$ induces on $\mathcal{C}(V, V^*)$ the structure of a braided rigid category if and only if $\mathcal{R}$ is dualisable.*

Proof. We have already explained one direction: suppose that $\mathcal{R}$ extends to a braiding Ψ to give the structure of a rigid braided monoidal category. Especially, it extends to a braiding between V and V^* which commutes with the adjunctions. One deduces easily that $\mathcal{R}$ is dualisable.

On the other hand, suppose that $\mathcal{R}$ is dualisable. By definition, we already have the braiding between V and V^* having the desired properties. Next, one can write out the braid relations or mixed quantum-Yang-Baxter equations on the space $V \otimes V^* \otimes V$ and applying $\text{ev}_V, \text{coev}_V$ one can deduce that v^{-1} exists. We can then define the braiding between V and V^{**} by identifying V and V^{**} via this map. In a similar way one extends inductively the braiding to ‘higher’ duals using the ‘ $\mathbb{Z}_2$ -periodicity’ $V^{[i+2]} \cong V^{[i]}$ induced by $v_{V^{[i]}}$. Formally, this means extending $\mathcal{R}$ using the formulas of Proposition 5.4. ■

Summing up the above remarks, we see that we can take the existence of a braided rigid category generated by V as an intrinsic definition of the dualisability. With some more work [12] one obtains from this an Abelian category, and can also improve on the requirements for dualisability. Another important construction in this context is that of quantum or categorical dimension. Thus in any rigid braided monoidal category there is an intrinsic categorical trace of any morphism $f : V \rightarrow V$ cf [22]

$$\text{Trace}_{\mathcal{R}}(f) := \text{ev}_V \circ \Psi_{V^*, V} \circ (\text{id} \otimes f) \circ \text{coev}_V.$$

The *categorical dimension* of any object is defined as the trace of the identity morphism. The use of this categorical construction to understand the so-called quantum dimensions in the theory of quantum groups was introduced in [19] and elsewhere. In our present setting the objects are vector spaces and the braiding is given by a dualisable Yang-Baxter operator $\mathcal{R}$, in which case the categorical dimension can also be thought of as a kind of trace of $\mathcal{R}$,

$$\dim_{\mathcal{R}}(V) := \text{Trace}(\mathcal{R}) := \text{ev} \circ \mathcal{R}^e \circ \text{coev}.$$

Proposition 5.6. *Let $(X, \mathcal{R})$ and $(Y, \mathcal{R}')$ be two dualisable q -Hecke Yang-Baxter operators. Then the maps $u_{X \oplus Y}$ and $v_{X \oplus Y}$ related with the glueing $\mathcal{R} \oplus_q \mathcal{R}'$ satisfy*

$$\begin{aligned} u_{X \oplus Y}(x) &= u_X(x), & u_{X \oplus Y}(y) &= (1 - (q - q^{-1}) \cdot \text{Trace}(\mathcal{R})) \cdot u_Y(y), \\ v_{X \oplus Y}(x) &= (1 - (q - q^{-1}) \cdot \text{Trace}(\mathcal{R}')) \cdot v_X(x), & v_{X \oplus Y}(y) &= v_Y(y), \end{aligned}$$

for $x \in X, y \in Y$. Moreover, the braided trace of the glueing can be computed as

$$\text{Trace}(\mathcal{R} \oplus_q \mathcal{R}') = \text{Trace}(\mathcal{R}) + \text{Trace}(\mathcal{R}') - (q - q^{-1}) \cdot \text{Trace}(\mathcal{R}) \cdot \text{Trace}(\mathcal{R}').$$

Proof. The proof is based on a direct verification. For example, we have

$$\begin{aligned}
u_{X \oplus Y}(x) &= (\text{ev}_{X \oplus Y} \otimes \text{id})(\Psi_{(X \oplus Y), (X \oplus Y)^*} \otimes \text{id})(\text{id} \otimes \text{coev}_{(X \oplus Y)^*})(x) \\
&= (\text{ev}_{X \oplus Y} \otimes \text{id})(\Psi_{(X \oplus Y), (X \oplus Y)^*} \otimes \text{id})(x \otimes (\text{coev}_{X^*} \oplus \text{coev}_{Y^*})) \\
&= (\text{ev}_{X \oplus Y} \otimes \text{id})(\Psi_{X, X^*} \otimes \text{id})(x \otimes \text{coev}_{X^*}) + (\text{ev}_{X \oplus Y} \otimes \text{id})(\Psi_{X, Y^*} \otimes \text{id})(x \otimes \text{coev}_{Y^*}) \\
&= u_X(x).
\end{aligned}$$

Here we used the fact that $\Psi_{(X \oplus Y), (X \oplus Y)^*}|_{X \otimes X^*} = \Psi_{X, X^*}$ and the fact that $\Psi_{(X \oplus Y), (X \oplus Y)^*}$ acts on $X \otimes Y^*$ by interchanging the arguments, which implies that $(\text{ev}_{X \oplus Y} \otimes \text{id})(\Psi_{X, Y^*} \otimes \text{id})(x \otimes \text{coev}_{Y^*}) = 0$. The verification of the remaining equations of the proposition is similar. $\blacksquare$

Example 5.7. Let $\mathcal{R}_{su(n)}$ be the classical Yang-Baxter operator introduced in Example 2.5. Clearly $\text{Trace}(\mathcal{R}_{su(1)}) = q^{-1}$ and the proposition above gives by induction the formula

$$\text{Trace}(\mathcal{R}_{su(n)}) = \sum_{i=1}^n q^{-2i+1} = \frac{1 - q^{-2n}}{1 - q^{-2}} q^{-1}.$$

A similar computation shows that $\text{Trace}(\mathcal{R}_{gl(1|1)})$ of the non-standard Yang-Baxter operator of Example 2.6 is zero. These are essentially the standard results of the quantum dimension of the fundamental representation of the associated quantum groups.

Example 5.8. For $V = \text{Span}(x_1, \dots, x_n)$ let $(x_1^{**}, \dots, x_n^{**})$ be the basis of V^{**} given by $x_i^{**}(\varphi) = \varphi(x_i)$, $\varphi \in V^*$. Using inductively Proposition 5.6 and the formula for $\text{Trace}(\mathcal{R}_{su(n)})$ computed in the previous example, we get the following formulas for the maps u_V and v_V related with $\mathcal{R}_{su(n)}$:

$$u_V(x_i) = q^{-2i+1} \cdot x_i^{**}, \quad v_V(x_i^{**}) = q^{-2n+2i-1} \cdot x_i,$$

for $1 \leq i \leq n$.

We close this section by discussing the following notion related with invariants of oriented framed isotopy classes of links. Let $\mathcal{R}$ be a dualisable Yang-Baxter operator on V and denote, as usual, by $\mathcal{R}^e$ its extension on $V \oplus V^*$. By the *double twist* we mean the map $\chi_{\mathcal{R}} : V \rightarrow V$ defined by

$$\chi_{\mathcal{R}} := (\text{id} \otimes \text{ev})(\mathcal{R}^e \otimes \text{id})(\mathcal{R}^e \otimes \text{id})(\text{coev} \otimes \text{id}) = (\text{id} \otimes \text{ev})(\mathcal{R}^{-1\#} \otimes \text{id})(\mathcal{R}^{\#-1} \otimes \text{id})(\text{coev} \otimes \text{id}).$$

One says that $\mathcal{R}$ is *tortile* [9] or *ribbon* [21] if there exists a map $\theta : V \rightarrow V$ (a *ribbon element*) such that $\chi_{\mathcal{R}} = \theta^2$. The following proposition shows that the structure of the double twist of q -Hecke Yang-Baxter operators is very simple.

Proposition 5.9. *Let $\mathcal{R}$ be a dualisable q -Hecke Yang-Baxter operator. Then*

$$\chi_{\mathcal{R}} = (1 - (q - q^{-1})\text{Trace}(\mathcal{R})) \cdot \mathbf{1}.$$

Proof. The q -Hecke property can be rewritten as $\mathcal{R}^{-1} = \mathcal{R} - (q - q^{-1}) \cdot (\mathbf{1} \otimes \mathbf{1})$. We have from this $\mathcal{R}^{-1\#} \circ \mathcal{R}^{\#-1} = (\mathbf{1} \otimes \mathbf{1}) - (q - q^{-1}) \cdot (\mathbf{1} \otimes \mathbf{1})^{\#} \circ (\mathcal{R}^{\#})^{-1}$. Plugging it into the formula defining $\chi_{\mathcal{R}}$ we get the requisite result. $\blacksquare$

Using Proposition 5.6 we easily get the following corollary.

Corollary 5.10. *Let $(X, \mathcal{R})$ and $(Y, \mathcal{R}')$ be two q -Hecke dualisable Yang-Baxter operators. Then*

$$\chi_{\mathcal{R} \oplus_q \mathcal{R}'} = \left[(1 - (q - q^{-1}) \cdot \text{Trace}(\mathcal{R})) \cdot (1 - (q - q^{-1}) \cdot \text{Trace}(\mathcal{R}')) \right] \cdot \mathbf{1}.$$

Thus, the scalar multiples by which the operators $\chi_{\mathcal{R}}$ act behave multiplicatively under $\oplus_q$.

As an immediate consequence of Proposition 5.9 we get that a dualisable q -Hecke Yang-Baxter operator $\mathcal{R}$ has a twist if and only if the scalar $(1 - (q - q^{-1}) \cdot \text{Trace}(\mathcal{R}))$ has a square root. Let $\lambda_{\mathcal{R}}$ be such a square root. Then the twist θ is necessarily of the form $\theta = \lambda_{\mathcal{R}} \cdot E$, where E is an endomorphism such that $E^2 = \mathbf{1}$. Combining the results above, we obtain the following proposition.

Proposition 5.11. *Let $(\mathcal{R}, \theta)$ and $(\mathcal{R}', \theta')$ be two tortile q -Hecke Yang-Baxter operators. Suppose $\theta = \lambda \cdot E$, $\theta' = \lambda' \cdot E'$, for some endomorphisms E and E' with $E^2 = \mathbf{1}$ and $E'^2 = \mathbf{1}$. If we put $\bar{\theta} := \lambda \cdot \lambda' \cdot (E \oplus E')$, then $(\mathcal{R} \oplus_q \mathcal{R}', \bar{\theta})$ is also a tortile Yang-Baxter operator.*

Example 5.12. From Example 5.7 we get easily

$$\chi_{\mathcal{R}_{su(n)}} = q^{-2n} \cdot \mathbf{1}, \quad \chi_{\mathcal{R}_{gl(1|1)}} = \mathbf{1}$$

(the formula for $\chi_{\mathcal{R}_{su(n)}}$ can be also obtained inductively from Corollary 5.10). Both operators are tortile and we usually take $\theta = q^{-n} \cdot \mathbf{1}$ for the first one and $\theta = \mathbf{1}$ for the second one. These are essentially the standard results for these quantum groups, cf [9][21].

6. Glueing of link invariants.

As we have already mentioned, there is an intimate relationship between link invariants and solutions of the Yang-Baxter equation. This suggest the possibility of the existence of an analog of glueing also for link invariants. The aim of this last section is to describe such a construction, based on the state-model approach of L. Kauffman [10].

For an (oriented) link, choose a plane projection without triple points and with only finitely many double points. We call such a projection the *shadow* of L . The isotopy type of L is then determined by its shadow plus the over/under information at the vertices (the double points) of the shadow. By a *state* of L we mean an assignment of an element of the set $\{x, y\}$ to each edge of the shadow. We will consider only states satisfying the *spin-conservation rule* meaning that at each double point of the shadow S ,

$$\begin{array}{ccc} i & & k \\ & \searrow & \nearrow \\ & \bullet & \\ & \nearrow & \searrow \\ j & & l \end{array} \quad i, j, k, l \in \{x, y\},$$

the equation $s(i) + s(k) = s(j) + s(l)$, with s defined by $s(x) = -1/2$, $s(y) = +1/2$, is satisfied.


Let L be a link and σ a state as above. It is convenient to classify the double points of the shadow and the related over/under information in a given state into the following types:

Type I: $\begin{array}{ccc} x & & x \\ & \searrow & \nearrow \\ & \bullet & \\ & \nearrow & \searrow \\ x & & x \end{array} \quad \text{and} \quad \begin{array}{ccc} y & & y \\ & \searrow & \nearrow \\ & \bullet & \\ & \nearrow & \searrow \\ y & & y \end{array},$

Type II: $\begin{array}{ccc} x & & x \\ & \nearrow & \searrow \\ & \bullet & \\ & \searrow & \nearrow \\ x & & x \end{array} \quad \text{and} \quad \begin{array}{ccc} y & & y \\ & \nearrow & \searrow \\ & \bullet & \\ & \searrow & \nearrow \\ y & & y \end{array},$

Type III: $\begin{array}{ccc} x & & y \\ & \searrow & \nearrow \\ & \bullet & \\ & \nearrow & \searrow \\ x & & y \end{array} \quad \text{and} \quad \begin{array}{ccc} y & & x \\ & \searrow & \nearrow \\ & \bullet & \\ & \nearrow & \searrow \\ y & & x \end{array},$


Type IV:



Type V:







Type VI:



By our spin-conservation rule, listed above are all possible situations.

Let L^σ be the singular (= possibly with double points) oriented link and state on it constructed from L and σ by the replacements:





the other types of vertices remaining unchanged.

Let L_y^σ be the component of L^σ colored by x and let L_x^σ have the similar meaning for coloring by y . Notice that the intersection $L_x^\sigma \cap L_y^\sigma$ need not be void, but L_x^σ and L_y^σ are honest oriented links (with no self-intersections).

Denote by V the set of vertices of the shadow S . Using our above classification of the types of vertices relative to our state, we define the *weight* $W^\sigma(v)$ of $v \in V$ by

$$W^\sigma(v) = \begin{cases} 1, & \text{for types I, II and VI,} \\ 0, & \text{for type III,} \\ (q - q^{-1}), & \text{for type IV,} \\ -(q - q^{-1}), & \text{for type V.} \end{cases}$$

Let L be an oriented link. Making the replacements



in the shadow S of L , we get a finite set of oriented circuits in the plane. Define $n(L) :=$ number of anticlockwise oriented circuits $-$ number of clockwise oriented circuits.


Consider a q -Hecke tortile Yang-Baxter operator $(\mathcal{R}, \theta)$. It is well-known (see, for example, [9]) that $(\mathcal{R}, \theta)$ gives rise to a framed oriented link isotopy invariant, say $\kappa(-)$. The idea of the construction of κ is to interpret a given link as a map in a certain free tortile category, the interpretation being based on the assignments

$$\begin{aligned} \text{Diagram 1} &\mapsto \mathcal{R}, & \text{Diagram 2} &\mapsto \mathcal{R}^{-1}, & \text{Diagram 3} &\mapsto \text{coev}, \\ \text{Diagram 4} &\mapsto \text{ev}, & \text{Diagram 5} &\mapsto (\mathbf{1} \otimes \theta^{-1}) \circ \mathcal{R}^e \circ \text{coev}, & \text{Diagram 6} &\mapsto \text{ev} \circ \mathcal{R}^e \circ (\mathbf{1} \otimes \theta^{-1}). \end{aligned}$$

Denote

$$\vartheta := \text{Cup} \downarrow, \quad \vartheta^{-1} := \text{Cap} \downarrow.$$

and let $\boxed{n}$ be the n -fold composition of ϑ for $n \geq 0$ and the $(-n)$ -fold composition of ϑ^{-1} for $n < 0$. Let, finally, Γ_n be the link



We can easily prove using the results of Section 5, the following proposition.

Proposition 6.1. *Let $(\mathcal{R}, \theta)$ and $(\mathcal{R}', \theta')$ be two q -Hecke tortile Yang-Baxter operators and suppose that $\theta = \lambda \cdot \mathbf{1}$ and $\theta' = \lambda' \cdot \mathbf{1}$ for some scalars λ and λ' . Let κ, κ' and ω be the oriented framed link isotopy invariants related with the operators $\mathcal{R}, \mathcal{R}'$ and $\mathcal{R} \oplus_q \mathcal{R}'$, respectively. Then*

$$\omega(\Gamma_n) = \lambda^{n-1} \kappa(\Gamma_n) + \lambda^{n+1} \kappa'(\Gamma_n).$$

Proof. From the assumption $\theta = \lambda \cdot \mathbf{1}$ we easily have that $\kappa(\Gamma_n) = \lambda^n \kappa(\Gamma_0)$ (notice that Γ_0 is the oriented circle). Immediately from the definition we get that $\kappa(\Gamma_0) = \text{ev} \circ \mathcal{R}^e \circ (\mathbf{1} \otimes \theta^{-1}) \circ \text{coev} = \lambda^{-1} \cdot \text{Trace}(\mathcal{R})$. This, and a similar computation gives

$$\kappa(\Gamma_n) = \lambda^{n-1} \cdot \text{Trace}(\mathcal{R}), \quad \kappa(\Gamma'_n) = \lambda'^{n-1} \cdot \text{Trace}(\mathcal{R}').$$

Recall that we consider the glueing $\mathcal{R} \oplus_q \mathcal{R}'$ as a tortile Yang-Baxter operator with the twist given by $\bar{\theta} = \lambda \lambda' \cdot \mathbf{1}$. The similar considerations as above together with Proposition 5.6 give that

$$\begin{aligned} \omega(\Gamma_n) &= \lambda^{n-1} \lambda'^{n-1} \cdot \text{Trace}(\mathcal{R} \oplus_q \mathcal{R}') \\ &= \lambda^{n-1} \lambda'^{n-1} \cdot [\text{Trace}(\mathcal{R}) + \text{Trace}(\mathcal{R}') - (q - q^{-1}) \cdot \text{Trace}(\mathcal{R}) \text{Trace}(\mathcal{R}')] = \\ &= \lambda^{n-1} [\lambda^{n-1} \text{Trace}(\mathcal{R})] + \lambda^{n-1} [\lambda'^{n-1} \text{Trace}(\mathcal{R}')] [1 - (q - q^{-1}) \cdot \text{Trace}(\mathcal{R})] \end{aligned}$$

which, together with the formula $\lambda^2 = 1 - (q - q^{-1}) \cdot \text{Trace}(\mathcal{R})$ (see the comments following Corollary 5.10) gives the requisite equation. ■

Let $(\mathcal{R}, \theta)$ and $(\mathcal{R}', \theta')$ be as in Proposition 6.1 and let κ and κ' be the related invariants.

Theorem 6.2. *Let κ, κ' be invariants of framed oriented isotopy and extend the definition by $\kappa(\emptyset) = \kappa'(\emptyset) := 1$ for the empty link $\emptyset$. Suppose, moreover, that*

$$\kappa(\vartheta) = \lambda \cdot \kappa(\downarrow), \quad \kappa(\vartheta^{-1}) = \lambda^{-1} \cdot \kappa(\downarrow), \quad \kappa'(\vartheta) = \lambda' \cdot \kappa'(\downarrow), \quad \kappa'(\vartheta^{-1}) = \lambda'^{-1} \cdot \kappa'(\downarrow),$$

for $\lambda, \lambda' \in \mathbf{k}$. Then their glueing $\kappa \oplus_q \kappa'$ defined by

$$(\kappa \oplus_q \kappa')(L) := \sum_{\text{states } \sigma} \kappa(L_x^\sigma) \kappa'(L_y^\sigma) \left(\prod_{v \in V} W^\sigma(v) \right) \lambda'^{-n(L_x^\sigma)} \lambda^{n(L_y^\sigma)}$$

is an invariant of framed oriented isotopy. Moreover, if κ, κ' correspond to q -Hecke Yang-Baxter operators as in Proposition 6.1 then their glueing $\kappa \oplus_q \kappa'$ corresponds to $\mathcal{R} \oplus_q \mathcal{R}'$.

Proof. We shall prove first that the stated formula for $\kappa \oplus_q \kappa'$ really defines a framed oriented isotopy invariant, which is the same as to prove that it is invariant under oriented framed Reidemeister moves [23, page 330]. This can be done by a direct verification, using the invariance of κ and κ' and the assumed values on $\vartheta, \vartheta^{-1}$. These values are the ones for the invariants corresponding to the Yang-Baxter operators in Proposition 6.1 where $\theta = \lambda \cdot \mathbf{1}$ and $\theta' = \lambda' \cdot \mathbf{1}$. It remains to prove that $\kappa \oplus_q \kappa'$ is related with the glueing $\mathcal{R} \oplus_q \mathcal{R}'$, in other words, we shall prove that $\omega = \kappa \oplus_q \kappa'$, where ω has the same meaning as in Proposition 6.1.

Observe that both oriented framed link isotopy invariants ω and $\kappa \oplus_q \kappa'$ satisfy the ‘skein relations’ in the form of [19, page 23] – for ω it follows from the fact that $\mathcal{R} \oplus_q \mathcal{R}'$ is q -Hecke while for $\kappa \oplus_q \kappa'$ it follows from the definition and from the fact that κ and κ' share this property, since $\mathcal{R}$ and $\mathcal{R}'$ are q -Hecke. Invoking the ‘skein template algorithm’ of [10, pages 57–63] we see that it is enough to verify that $\omega(\Gamma_n) = (\kappa \oplus_q \kappa')(\Gamma_n)$ for all n , but the left hand side was already computed in of Proposition 6.1 and it is easy to see that it coincides with the value of $\kappa \oplus_q \kappa'$ from its definition. ■

Example 6.3. Let us recall first the definition of the *writhe* $w(L)$ of an oriented link L . We put

$$w(L) = \sum_{v \in V} \epsilon(v),$$

where the summation is taken over the set of vertices of the shadow of L and ϵ is defined by

$$\epsilon \left(\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \right) := +1, \quad \epsilon \left(\begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} \right) := -1.$$

Consider now $\mathcal{R}_{su(1)}$ as a tortile Yang-Baxter operator with $\theta = q \cdot \mathbb{1}$. It is easy to show that the related invariant is given by $\kappa(L) = q^{w(L)}$. We show that the formula in Theorem 6.2 for the glueing $\kappa \oplus_q \kappa$ gives the usual state model for the $\mathcal{R}_{su(2)}$ -Jones invariant, as predicted by Theorem 6.2. We have

$$(17) \quad (\kappa \oplus_q \kappa)(L) := \sum_{\sigma} q^{w(L_x^{\sigma}) + w(L_y^{\sigma})} \left(\prod_{v \in V} W^{\sigma}(v) \right) q^{-n(L_x^{\sigma})} q^{n(L_y^{\sigma})}.$$

Recall our classification of the types of vertices of the shadow of L and define, for each vertex $v \in V$,

$$U^{\sigma}(v) := \begin{cases} q, & \text{for type I,} \\ q^{-1}, & \text{for type II,} \\ 1, & \text{for types III–VI.} \end{cases}$$

It is immediate from the definition that

$$q^{w(L_x^{\sigma}) + w(L_y^{\sigma})} = \prod_{v \in V} U^{\sigma}(v).$$

Looking at $W_{\mathcal{R}_{su(2)}}^{\sigma}(v) := U^{\sigma}(v) \cdot W^{\sigma}(v)$, we see that $W_{\mathcal{R}_{su(2)}}^{\sigma}(v)$ is the usual Boltzmann weight given by the formula

$$W_{\mathcal{R}_{su(2)}}^{\sigma}(v) = \begin{cases} q, & \text{for type I,} \\ q^{-1}, & \text{for type II,} \\ 0, & \text{for type III,} \\ (q - q^{-1}), & \text{for type IV,} \\ -(q - q^{-1}), & \text{for type V,} \\ 1, & \text{for type VI} \end{cases}$$

Summing up the above observations we see that the formula (17) is identical with Kauffman's state-model definition of the $\mathcal{R}_{su(2)}$ -Jones invariant. We used the conventions in [19, par. 2.2].

References.

- [1] P. Deligne. Catégories Tannakiennes. *Preprint*, 1988.
- [2] V.G. Drinfeld. Quantum groups. In A. Gleason, editor, *Proceedings of the ICM*, pages 798–820, Rhode Island, 1987. AMS.
- [3] L.D. Faddeev, N.Yu. Reshetikhin, and L.A. Takhtajan. Quantization of Lie groups and Lie algebras. *Leningrad Math J.*, 1:193–225, 1990.
- [4] P. Freyd and D. Yetter. Braided compact closed categories with applications to low dimensional topology. *Adv. Math.*, 77:156–182, 1989.
- [5] M-L. Ge, N. Jing, and Y-S. Wu. A new quantum group associated with “non-standard” braid group representation. *Preprint*, 1990.

- [6] D.I. Gurevich. Algebraic aspects of the quantum Yang-Baxter equation. *Leningrad Math. J.*, 2:801–828, 1991.
- [7] P.J. Hilton and U.S. Stammbach. *A Course in Homological Algebra*. Springer-Verlag, 1980.
- [8] A. Joyal and R. Street. Braided monoidal categories. Mathematics Reports 86008, Macquarie University, 1986.
- [9] A. Joyal and R. Street. Tortile Yang-Baxter operators in tensor categories. *J. Pure Applied Algebra*, 71:43–51, 1991.
- [10] L. Kauffman. *Knots and Physics*, volume 1 of *Knots and Everything*. World Sci., 1991.
- [11] V.V. Lyubashenko. Hopf algebras and vector symmetries. *Russ. Math. Surveys*, 41:153–154, 1986.
- [12] V.V. Lyubashenko. *Superanalysis and solutions to the triangles equation*, PhD thesis, Kiev 1986.
- [13] S. Majid. Algebras and Hopf algebras in braided categories. In *Advances in Hopf Algebras*. In press, Plenum.
- [14] S. Majid. Anyonic quantum groups. In Z. Oziewicz et al, eds, *Spinors, twistors, Clifford algebras and quantum deformations (Proc. of 2nd Max Born Symposium, Wroclaw, Poland, 1992)*, pages 327–336. Kluwer.
- [15] S. Majid. Braided momentum in the q -Poincaré group. *J. Math. Phys.*, 34:2045–2058, 1993.
- [16] S. Majid. Examples of braided groups and braided matrices. *J. Math. Phys.*, 32:3246–3253, 1991.
- [17] S. Majid. Free braided differential calculus, braided binomial theorem and the braided exponential map. In press in *J. Math. Phys.* 34, 1993.
- [18] S. Majid. More examples of bicrossproduct and double cross product Hopf algebras. *Isr. J. Math*, 72:133–148, 1990.
- [19] S. Majid. Quasitriangular Hopf algebras and Yang-Baxter equations. *Int. J. Modern Physics A*, 5(1):1–91, 1990.
- [20] S. Majid. Quantum and braided linear algebra. *J. Math. Phys.*, 34:1176–1196, 1993.
- [21] N.Yu. Reshetikhin and V.G. Turaev. Ribbon graphs and their invariants derived from quantum groups. *Comm. Math. Phys.*, 127(1):1–26, 1990.
- [22] N. Saavedra Rivano. Catégories Tannakiennes. *Springer Lect. Notes in Math*, 265, 1972.
- [23] D.N. Yetter. Framed tangles and a theorem of Deligne on braided deformations of Tannakian categories. *Contemp. Mathematics*, 134:325–349, 1992.